

Store at
-20C
#39898**iNOS Antibody**

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 130	Source/Isotype: Rabbit	UniProt ID: #P35228	Entrez-Gene Id: 4843
-------------------------------	-------------------------	-----------------------------------	-------------------------	----------------------------------	-------------------------------	--------------------------------

Product Usage Information**Application**

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. *Do not aliquot the antibody.*

Specificity/Sensitivity

iNOS Antibody recognizes endogenous levels of total iNOS protein.

Source / Purification

Polyclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human iNOS protein. Antibodies are purified by peptide affinity chromatography.

Background

Nitric Oxide Synthase (NOS) catalyzes the formation of nitric oxide (NO) and citrulline from L-arginine, oxygen, and cofactors. Three family members have been characterized: neuronal NOS (nNOS), which is found primarily in neuronal tissue; inducible NOS (iNOS), which is induced by interferon gamma and lipopolysaccharides in the kidney and cardiovascular system; and endothelial NOS (eNOS), which is expressed in blood vessels (1). NO is a messenger molecule with diverse functions throughout the body, including the maintenance of vascular integrity, homeostasis, synaptic plasticity, long-term potentiation, learning, and memory (2,3).

NO catalyzed by iNOS is involved in host defense against protozoa, bacteria, fungi, and viruses. Unlike constitutively expressed eNOS and nNos, iNOS is not usually expressed in quiescent cells. iNOS is transcriptionally induced in response to bacterial endotoxins, such as LPS and proinflammatory cytokines, in macrophages and various other cell types. Transcription factors involved in iNOS transcription include NF-κB, AP-1, and STAT. Different signaling pathways either promote (Jak1/2, PKC, c-Raf, p38 MAP kinase, and p44/42 MAP kinase) or inhibit (PI3 kinase) iNOS expression depending on stimulus and cell type (4).

Background References

1. Tsutsui, M. (2004) *J Atheroscler Thromb* 11, 41-8.
2. Son, H. et al. (1996) *Cell* 87, 1015-23.
3. Hawkins, R.D. (1996) *Neuron* 16, 465-7.
4. Bogdan, C. (2001) *Nat Immunol* 2, 907-16.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.