

| | | |
|--|---|---|
| #4022 Store at -20C | MMP-2 Antibody |  |
| | Orders: 877-616-CELL (2355) orders@cellsignal.com | |
| | Support: 877-678-TECH (8324) | |
| | Web: info@cellsignal.com cellsignal.com | |
| 3 Trask Lane Danvers Massachusetts 01923 USA | | |

For Research Use Only. Not for Use in Diagnostic Procedures.

| Applications: | Reactivity: | Sensitivity: | MW (kDa): | Source/Isotype: | UniProt ID: | Entrez-Gene Id: |
|---------------|-------------|--------------|-----------|-----------------|-------------|-----------------|
| W | H | Endogenous | 64, 72 | Rabbit | #P08253 | 4313 |

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

MMP-2 Antibody detects full length (proenzyme, 72 kDa) and cleaved (active enzyme, 64 kDa) MMP-2.

Species predicted to react based on 100% sequence homology

Mouse, Rat

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Lys116 of human MMP-2. Antibodies are purified by protein A and peptide affinity chromatography.

Background

The matrix metalloproteinases (MMPs) are a family of proteases that target many extracellular proteins including other proteases, growth factors, cell surface receptors, and adhesion molecules (1). Among the family members, MMP-2, MMP-3, MMP-7, and MMP-9 have been characterized as important factors for normal tissue remodeling during embryonic development, wound healing, tumor invasion, angiogenesis, carcinogenesis, and apoptosis (2-4). Research studies have shown that MMP activity correlates with cancer development (2). One mechanism of MMP regulation is transcriptional (5). Once synthesized, MMP exists as a latent proenzyme. Maximum MMP activity requires proteolytic cleavage to generate active MMPs by releasing the inhibitory propeptide domain from the full-length protein (5).

Background References

- McCawley, L.J. and Matrisian, L.M. (2001) *Curr Opin Cell Biol* 13, 534-40.
- Coussens, L.M. et al. (2002) *Science* 295, 2387-92.
- Sternlicht, M.D. et al. (1999) *Cell* 98, 137-46.
- Vu, T.H. et al. (1998) *Cell* 93, 411-22.
- Nagase, H. et al. (1990) *Biochemistry* 29, 5783-9.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for

diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.