

Phospho-Stat3 (Tyr705) (D3A7) XP[®] Rabbit mAb (Biotinylated)**Orders:** 877-616-CELL (2355)
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R Mk	Endogenous	79, 86	Rabbit IgG	#P40763	6774

Product Usage Information	Application	Dilution
	Western Blotting	1:1000
Storage	Supplied in 136 mM NaCl, 2.6 mM KCl, 12 mM sodium phosphate (pH 7.4) dibasic, 2 mg/ml BSA, and 50% glycerol. Store at -20°C. Do not aliquot the antibodies.	
Specificity/Sensitivity	Phospho-Stat3 (Tyr705) (D3A7) XP [®] Rabbit mAb (Biotinylated) detects endogenous levels of Stat3 only when phosphorylated at Tyr705. This antibody does not cross-react with phospho-EGFR or the corresponding phospho-tyrosines of other Stat proteins.	
Species predicted to react based on 100% sequence homology	Hamster, Bovine, Pig, Horse	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr705 of mouse Stat3.	
Description	This Cell Signaling Technology (CST) antibody is conjugated to biotin under optimal conditions. The unconjugated Phospho-Stat3 (Tyr705) (D3A7) XP [®] Rabbit mAb #9145 reacts with human, mouse, rat and monkey phospho-Stat3 (Tyr705) protein. CST expects that Phospho-Stat3 (Tyr705) (D3A7) XP [®] Rabbit mAb (Biotinylated) will also recognize phospho-Stat3 (Tyr705) in these species.	
Background	The Stat3 transcription factor is an important signaling molecule for many cytokines and growth factor receptors (1) and is required for murine fetal development (2). Research studies have shown that Stat3 is constitutively activated in a number of human tumors (3,4) and possesses oncogenic potential (5) and anti-apoptotic activities (3). Stat3 is activated by phosphorylation at Tyr705, which induces dimerization, nuclear translocation, and DNA binding (6,7). Transcriptional activation seems to be regulated by phosphorylation at Ser727 through the MAPK or mTOR pathways (8,9). Stat3 isoform expression appears to reflect biological function as the relative expression levels of Stat3 α (86 kDa) and Stat3 β (79 kDa) depend on cell type, ligand exposure, or cell maturation stage (10). It is notable that Stat3 β lacks the serine phosphorylation site within the carboxy-terminal transcriptional activation domain (8).	
Background References	<ol style="list-style-type: none"> 1. Heim, M.H. (2001) <i>J Recept Signal Transduct Res</i> 19, 75-120. 2. Takeda, K. et al. (1997) <i>Proc Natl Acad Sci U S A</i> 94, 3801-4. 3. Catlett-Falcone, R. et al. (1999) <i>Immunity</i> 10, 105-15. 4. Garcia, R. and Jove, R. (1998) <i>J Biomed Sci</i> 5, 79-85. 5. Bromberg, J.F. et al. (1999) <i>Cell</i> 98, 295-303. 6. Darnell, J.E. et al. (1994) <i>Science</i> 264, 1415-21. 7. Ihle, J.N. (1995) <i>Nature</i> 377, 591-4. 8. Wen, Z. et al. (1995) <i>Cell</i> 82, 241-50. 9. Yokogami, K. et al. (2000) <i>Curr Biol</i> 10, 47-50. 10. Biethahn, S. et al. (1999) <i>Exp Hematol</i> 27, 885-94. 	

Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.
Applications Key	W: Western Blotting
Cross-Reactivity Key	H: Human M: Mouse R: Rat Mk: Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.