



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#4155

EPAC1 (5D3) Mouse mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 100	Source/Isotype: Mouse IgG2a	UniProt ID: #O95398	Entrez-Gene Id: 10411
---------------------------	-----------------------------	-----------------------------------	-------------------------	---------------------------------------	-------------------------------	---------------------------------

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

Epac1 (5D3) Mouse mAb detects endogenous levels of total Epac1 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a recombinant fragment of human Epac1.

Background

EPAC1 and EPAC2 (exchange proteins activated by cyclic AMP) are guanine nucleotide exchange factors (GEFs) that catalyze the exchange of GDP for GTP, activating Rap1 and Rap2 small GTPases. Rap activation by EPAC is cAMP-dependent and mediates cAMP signaling in part through protein kinase A (PKA) (reviewed in 1). EPAC signaling plays a significant role in a number of cellular processes including migration and focal adhesion formation (2), exocytosis (3), insulin signaling (4), axon growth and guidance (5) and neurotransmitter release (6).

Background References

1. Bos, J.L. (2006) *Trends Biochem Sci* 31, 680-6.
2. Lyle, K.S. et al. (2008) *Cell Signal* 20, 1104-16.
3. Branham, M.T. et al. (2009) *J Biol Chem* 284, 24825-39.
4. Petersen, R.K. et al. (2008) *Mol Cell Biol* 28, 3804-16.
5. Murray, A.J. and Shewan, D.A. (2008) *Mol Cell Neurosci* 38, 578-88.
6. Ouyang, M. et al. (2008) *Proc Natl Acad Sci USA* 105, 11993-7.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party,

whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.