Store at -20C

.209

-

## Phospho-NMDA Receptor 2B (GluN2B) (Tyr1070) Antibody



Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	<b>Reactivity:</b> R	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 200	<b>Source/Isotype:</b> Rabbit	<b>UniProt ID:</b> #Q13224	Entrez-Gene Id: 2904		
Product Usage Information		<b>Application</b> Western Blotting			Dilution 1:1000			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store 20°C. Do not aliquot the antibody.				ycerol. Store at –		
		Phospho-NMDA Receptor 2B (GluN2B) (Tyr1070) Antibody detects endogenous levels of NMDA Receptor 2B only when phosphorylated at Tyr1070.						
Species predict based on 100% homology		Human, Mouse						
Source / Purific	ation	Polyclonal antibodies are produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Tyr1070 of NMDA Receptor 2B. Antibodies are purified by protein A and peptide affinity chromatography.						
Background		N-methyl-D-aspartate receptor (NMDAR) forms a heterodimer of at least one NR1 and one NR2A-D subunit. Multiple receptor isoforms with distinct brain distributions and functional properties arise by selective splicing of the NR1 transcripts and differential expression of the NR2 subunits. NR1 subunits bind the co-agonist glycine and NR2 subunits bind the neurotransmitter glutamate. Activation of the NMDA receptor or opening of the ion channel allows flow of Na <sup>+</sup> and Ca <sup>2+</sup> ions into the cell, and K <sup>+</sup> out of the cell (1). Each subunit has a cytoplasmic domain that can be directly modified by the protein kinase/phosphatase (2). PKC can phosphorylate the NR1 subunit (NMDAR1) of the receptor at Ser890/Ser896, and PKA can phosphorylate NR1 at Ser897 (3). The phosphorylation of NR1 by PKC decreases its affinity for calmodulin, thus preventing the inhibitory effect of calcineurin on the receptor (5). NMDAR mediates long-term potentiation and slow postsynaptic excitation, which play central roles in learning, neurodevelopment, and neuroplasticity (6). Phosphorylation of NMDAR2B at Tyr1070 was identified at Cell Signaling Technology (CST) using PhosphoScan <sup>®</sup> , CST's MS/MS platform for phosphorylation site discovery. Phosphorylation of NMDAR2B at Tyr1070 was observed in extracts isolated from ischemic rat brain. For additional information please visit PhosphoSitePlus <sup>®</sup> , CST's modification site knowledgebase, at www.phosphosite.org.						
Background Re	ferences	1. Liu, X.B. et al. (2004) <i>J Neurosci</i> 24, 8885-95. 2. Westphal, R.S. et al. (1999) <i>Science</i> 285, 93-6. 3. Tingley, W.G. et al. (1997) <i>J Biol Chem</i> 272, 5157-66. 4. Hisatsune, C. et al. (1997) <i>J Biol Chem</i> 272, 20805-10. 5. Raman, I.M. et al. (1996) <i>Neuron</i> 16, 415-21. 6. Makhinson, M. et al. (1999) <i>J Neurosci</i> 19, 2500-10.						
Species Reactiv	vity	Species reactivity is de	termined by testing	g in at least one approve	ed application (e.g.,	western blot).		
Western Blot B	uffer		IT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X Tween® 20 at 4°C with gentle shaking, overnight.					
Applications Ke	ey	W: Western Blotting						
Cross-Reactivit	у Кеу	R: Rat						
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.						

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

## **Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party. whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.