

## SMYD2 Antibody



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H M R Mk	Endogenous	49	Rabbit	#Q9NRG4	56950

### Product Usage Information

#### Application

Western Blotting

#### Dilution

1:1000

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

### Specificity/Sensitivity

SMYD2 Antibody detects endogenous levels of total SMYD2 protein. The antibody does not cross-react with SMYD3 protein.

### Species predicted to react based on 100% sequence homology

Bovine, Pig

### Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to human SMYD2 protein. Antibodies are purified by protein A and peptide affinity chromatography.

### Background

SET and MYND domain-containing protein 2 (SMYD2), also known as lysine methyltransferase protein 3C (KMT3C), is a member of the SMYD family of protein methyltransferases (1). All five members of this family (SMYD1, SMYD2, SMYD3, SMYD4, and SMYD5) contain a conserved catalytic SET domain, originally identified in *Drosophila* Su[var]3-9, Enhancer of zeste, and Trithorax proteins. This domain is split by the MYN domain/zinc finger motif believed to facilitate protein-protein interactions (1). SMYD2 localizes to both the cytoplasm and nucleus, and is highly expressed in the adult mouse heart, brain, liver, kidney, thymus, and ovary, as well as in the developing mouse embryo (1). SMYD2 functions to repress transcription by interacting with the Sin3A repressor complex and methylating Lys36 of histone H3 (1). SMYD2 also interacts with HSP90α and methylates Lys4 of histone H3, a mark associated with transcriptional activation (2). In addition to histones as methyl substrates, SMYD2 methylates p53 at Lys370 to repress p53-mediated transcriptional activation and apoptosis (3,4).

### Background References

1. Brown, M.A. et al. (2006) *Mol Cancer* 5, 26.
2. Abu-Farha, M. et al. (2008) *Mol Cell Proteomics* 7, 560-72.
3. Huang, J. et al. (2006) *Nature* 444, 629-32.
4. Huang, J. et al. (2007) *Nature* 449, 105-8.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

### Applications Key

**W:** Western Blotting

### Cross-Reactivity Key

**H:** Human **M:** Mouse **R:** Rat **Mk:** Monkey

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

### Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.