ka hnRNP A1 (K350) Antibody





Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

_

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP, IF-IC	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 34, 39 kd	Source/Isotype: Rabbit	UniProt ID: #P09651	Entrez-Gene Id: 3178		
Product Usage Information	2	Application Western Blotting Immunoprecipitation Immunofluorescence	(Immunocytochem	istry)		Dilution 1:1000 1:50 1:50		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.						
Specificity/Sensitivity		hnRNP A1 (K350) Antibody detects both isoforms of endogenous hnRNP A1 protein.						
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Lys350 of human hnRNP A1 protein. Antibodies were purified by protein A and peptide affinity chromatography.						
Background		Heterogeneous nuclear ribonucleoprotein A1 (hnRNP A1) is a member of the hnRNP A/B family of related RNA binding proteins that bind pre-mRNA and are involved in the processing, metabolism, and transport of nuclear pre-mRNA transcripts (1). hnRNP A1 regulates the alternative splicing of c-Src and c-H-Ras (2,3) and modifies initiation of translation of the fibroblast growth factor 2 mRNA (4). hnRNP A1 expression level is elevated in many cancers; knockdown of hnRNP A1 leads to apoptosis in various cancer cells (5). Although predominantly nuclear, hnRNP A1 is continually transported from the nucleus to the cytoplasm where it disassociates from mRNA and is rapidly re-imported into the nucleus (6,7). hnRNP A1 binds to cis-acting repressive sequences (CRS) of HIV-1 to influence HIV-1 production (8,9). HIV-1 enhances hnRNP A1 expression and promotes the relocalization of hnRNP A1 to the cytoplasm (10).						
Background R	eferences	1. Myer, V.E. and Steitz 2. Rooke, N. et al. (200 3. Guil, S. et al. (2003) 4. Bonnal, S. et al. (2003 5. Patry, C. et al. (2003 6. Piñol-Roma, S. and J 7. Siomi, M.C. et al. (19 8. Black, A.C. et al. (19 9. Hadian, K. et al. (200 10. Monette, A. et al. (200	 Mol Cell Biol 23, Mol Cell Biol 23, 29 J Biol Chem 280, Cancer Res 63, 76 Dreyfuss, G. (1992) Cell Biol 138, Virus Genes 12, J Biol Chem 284 	1874-84. 27-41. 4144-53. 79-88. <i>Nature</i> 355, 730-2. 1181-92. 275-85. , 33384-91.				
Species Reacti	vity	Species reactivity is de	etermined by testin	g in at least one approve	ed application (e.g.,	western blot).		
Western Blot E	Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.						
Applications K	ey	W: Western Blotting II	P: Immunoprecipita	ation IF-IC: Immunofluor	rescence (Immunoc	ytochemistry)		
Cross-Reactivi	ty Key	H: Human M: Mouse R: Rat Mk: Monkey						
Trademarks ar	nd Patents	Cell Signaling Technol	ogy is a trademark	of Cell Signaling Techno	logy, Inc.			
		All other trademarks a more information.	are the property of	their respective owners.	Visit cellsignal.com	/trademarks for		
Limited Uses				a writing signed by a leg ovided by CST, its affiliate				

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.