

Anti-Rabbit IgG (H+L), F(ab')₂ Fragment (Alexa Fluor® 647 Conjugate)

✓ 250 µl



Orders ■ 877-616-CELL (2355)
orders@cellsignal.com

Support ■ 877-678-TECH (8324)
info@cellsignal.com

Web ■ www.cellsignal.com


rev. 09/19/17

For Research Use Only. Not For Use In Diagnostic Procedures.

Description: Anti-rabbit IgG (H+L) F(ab')₂ Fragment was conjugated to Alexa Fluor® 647 fluorescent dye under optimal conditions and formulated at 2 mg/ml. This F(ab')₂ fragment results in less non-specific binding to cells through Fc receptors.

Background: This product has been optimized for use as a secondary antibody in immunofluorescent applications. Fluorescent anti-species IgG conjugates are ideal for flow cytometry and immunofluorescence. Cell Signaling Technology's strict quality control procedures assure that each conjugate provides optimal specificity and fluorescence.

Specificity/Sensitivity: F(ab')₂ fragments are prepared from goat antibodies that have been adsorbed against pooled human serum, mouse serum, plasmacytoma/hybridoma proteins and purified human paraproteins.



Flow cytometric analysis of Jurkat cells, untreated (green) or treated with LY294002 #9901, wortmannin #9951 and U0126 #9906 (blue), using Phospho-S6 Ribosomal Protein (Ser235/236) (D57.2.2E) XP® Rabbit mAb #4858 detected with Anti-Rabbit IgG (H+L), F(ab')₂ Fragment (Alexa Fluor® 647 Conjugate) compared to a nonspecific negative control antibody (red).

Storage: Supplied in 0.1 M sodium phosphate, 0.1 M sodium chloride, pH 7.5, 5 mM sodium azide. Store at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.

Directions for Use: The optimal dilution of the anti-species antibody should be determined for each primary antibody by titration. However, a final dilution of 1:500 – 1:2000 should yield acceptable results for immunofluorescent and flow cytometry assays.





For product specific protocols and a complete listing of recommended companion products please see the product web page at www.cellsignal.com

This product is provided under an intellectual property license from Life Technologies Corporation. The transfer of this product is contingent on the buyer using the purchased product solely in research, including use with HCS or other automated imaging applications but excluding use in combination with DNA microarrays. The buyer must not sell or otherwise transfer this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; (c) manufacturing or quality assurance or quality control, or (d) resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com

Alexa Fluor is a registered trademark of Life Technologies Corporation.

DRAQ5 is a registered trademark of Biostatus Limited.

Acumen is a registered trademarks of TTP Labtech



High content analysis of A549 cells exposed to varying concentrations of LY294002 (#9901) for 3 hrs, followed by 100 ng/mL EGF for 20 minutes. With increasing concentrations of LY294002, a significant decrease (~5 fold) in phospho-S6 Ribosomal Protein (Ser235/236) signal as compared to the uninhibited control was observed. When using phospho-S6 as a measurement, the IC₅₀ of this compound was 3.06 µM. Data were generated on the Acumen® HCS platform using Anti-Rabbit IgG (H+L), F(ab')₂ Fragment (Alexa Fluor® 647 Conjugate).

◀ Confocal immunofluorescent analysis of mitotic HeLa cells using β-Tubulin (9F3) Rabbit mAb #2128 detected with Anti-Rabbit IgG (H+L), F(ab')₂ Fragment (Alexa Fluor® 647 Conjugate) (blue) and Phospho-Histone H3 (Ser10) (D2C8) XP® Rabbit mAb (Alexa Fluor® 555 Conjugate) #3475 (red).