

Caspase-5 Antibody



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H	Endogenous	48, 20	Rabbit	#P51878	838

Product Usage Information

Application

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

Caspase-5 Antibody detects endogenous levels of total caspase-5 protein. This antibody detects the p20 subunit of active caspase-5.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Leu130 within the p20 subunit of human caspase-5 protein. Antibodies were purified by protein A and peptide affinity chromatography.

Background

Caspase-5 (Ich-3/ICE_{rel}III/TY) is a member of the caspase family of cysteine proteases that play a key role in the execution of apoptosis and activation of inflammatory cytokines (1-3). Caspase-5 is widely expressed, with highest expression observed in placenta and lung (1). Interferon-γ and LPS regulate expression of caspase-5 (2,4). Members of the caspase-1 subfamily of caspases, which includes caspase-4, -5, and murine caspase-11 and -12, can induce apoptosis when over-expressed and mediate the proteolytic activation of inflammatory cytokines (5). Processing of IL-1β occurs through the activation of an inflammasome complex consisting of caspase-1, caspase-5, Pycard and NALP1 (6). Transcription factor Max, a component of the Myc/Mad/Max network, is cleaved by caspase-5 during Fas-induced apoptosis (7). Several alternative spliced variants of caspase-5 have been identified (8). Frameshift mutations of caspase-5 have been observed in leukemia, lymphoma (9), and colorectal cancers (10).

Background References

1. Munday, N.A. et al. (1995) *J Biol Chem* 270, 15870-6.
2. Wang, S. et al. (1996) *J Biol Chem* 271, 20580-7.
3. Faucheu, C. et al. (1996) *Eur J Biochem* 236, 207-13.
4. Lin, X.Y. et al. (2000) *J Biol Chem* 275, 39920-6.
5. Martinon, F. and Tschopp, J. (2007) *Cell Death Differ* 14, 10-22.
6. Martinon, F. et al. (2002) *Mol Cell* 10, 417-26.
7. Krippner-Heidenreich, A. et al. (2001) *Biochem J* 358, 705-15.
8. Eckhart, L. et al. (2006) *Biochem Biophys Res Commun* 348, 682-8.
9. Takeuchi, S. et al. (2003) *Leuk Res* 27, 359-61.
10. Trojan, J. et al. (2004) *Int J Colorectal Dis* 19, 538-44.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.