

Na,K-ATPase β 1 (D6U8Q) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene ID:
W	H M R	Endogenous	45-55	Rabbit IgG	#P05026	481

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C . Do not aliquot the antibody.

Specificity/Sensitivity

Na,K-ATPase β 1 (D6U8Q) Rabbit mAb recognizes endogenous levels of total Na,K-ATPase β 1 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro200 of human Na,K-ATPase β 1 protein.

Background

The Na,K-ATPase is an integral membrane heterodimer belonging to the P-type ATPase family. This ion channel uses the energy derived from ATP hydrolysis to maintain membrane potential by driving sodium export and potassium import across the plasma membrane against their electrochemical gradients. It is composed of a catalytic α subunit and a β subunit (reviewed in 1). Several phosphorylation sites have been identified for the α 1 subunit. Tyr10 is phosphorylated by an as yet undetermined kinase (2), Ser16 and Ser23 are phosphorylated by PKC, and Ser943 is phosphorylated by PKA (3-5). All of these sites have been implicated in the regulation of enzyme activity in response to hormones and neurotransmitters, altering trafficking and kinetic properties of Na,K-ATPase. Altered phosphorylation in response to angiotensin II stimulates activity in the rat proximal tubule (6). Na,K-ATPase is also involved in other signal transduction pathways. Insulin regulates its localization in differentiated primary human skeletal muscle cells, and this regulation is dependent on ERK1/2 phosphorylation of the α subunit (7). Na,K-ATPase and Src form a signaling receptor complex that affects regulation of Src kinase activity and, subsequently, its downstream effectors (8,9). Na,K-ATPase β 1 is the non-catalytic subunit of Na,K-ATPase. It is required for stabilization, maturation, and translocation of the catalytic α subunit to the plasma membrane (10-12). Na,K-ATPase β 1 also mediates the trans-dimerization of Na,K-ATPase between neighboring cells where it regulates the integrity of tight junctions (13-17). Glutathionylation of Na,K-ATPase β 1 regulates the ion pump activity of Na,K-ATPase (18). Research studies have shown that Na,K-ATPase β 1 is a target of the Sonic Hedgehog signaling pathway and may be involved in suppressing tumor development and progression (19). *ATP1B1*, the gene encoding Na,K-ATPase β 1, is epigenetically silenced by promoter methylation in both renal cell carcinoma cell lines and patient tissues (20).

Background References

1. Therien, A.G. and Blostein, R. (2000) *Am J Physiol Cell Physiol* 279, C541-66.
2. Féraille, E. et al. (1999) *Mol Biol Cell* 10, 2847-59.
3. Fisone, G. et al. (1994) *J Biol Chem* 269, 9368-73.
4. Feschenko, M.S. and Sweadner, K.J. (1995) *J Biol Chem* 270, 14072-7.
5. Beguin, P. et al. (1994) *J Biol Chem* 269, 24437-45.
6. Yingst, D.R. et al. (2004) *Am J Physiol Renal Physiol* 287, F713-21.
7. Al-Khalili, L. et al. (2004) *J Biol Chem* 279, 25211-8.
8. Tian, J. et al. (2006) *Mol Biol Cell* 17, 317-26.
9. Liang, M. et al. (2006) *J Biol Chem* 281, 19709-19.
10. Beggah, A.T. et al. (1997) *J Biol Chem* 272, 10318-26.
11. Hasler, U. et al. (1998) *J Biol Chem* 273, 30826-35.
12. Rajasekaran, S.A. et al. (2004) *Mol Biol Cell* 15, 3224-32.
13. Rajasekaran, S.A. et al. (2001) *Mol Biol Cell* 12, 3717-32.
14. Rajasekaran, A.K. and Rajasekaran, S.A. (2003) *Am J Physiol Renal Physiol* 285, F388-96.
15. Bab-Dinitz, E. et al. (2009) *Biochemistry* 48, 8684-91.
16. Tokhtaeva, E. et al. (2011) *J Biol Chem* 286, 25801-12.
17. Vagin, O. et al. (2012) *Am J Physiol Cell Physiol* 302, C1271-81.
18. Figtree, G.A. et al. (2012) *Free Radic Biol Med* 53, 2263-8.
19. Lee, S.J. et al. (2015) *Mol Cancer* 14, 159.
20. Selvakumar, P. et al. (2014) *Epigenetics* 9, 579-86.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.