

Rab27B Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 27	Source/Isotype: Rabbit	UniProt ID: #000194	Entrez-Gene Id: 5874
Product Usage Information		Application Western Blotting			Dilution 1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at – 20°C. <i>Do not aliquot the antibody.</i>				
Specificity/Sensitivity		Rab27B Antibody recognizes endogenous levels of total Rab27B protein. Overexpression analysis confirms that this antibody does not detect Rab27A.				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human Rab27B protein. Antibodies are purified by protein A and peptide affinity chromatography.				
Background		Rab27 proteins are members of the Ras superfamily of small Rab GTPases associated with the process of exocytosis (1-2), and in particular the exosome secretion pathway (3-5). There are two Rab27 isoforms, Rab27A and Rab27B, which are encoded by separate genes and which exhibit distinct subcellular localization and functions. Rab27A colocalizes to CD63 ⁺ vesicles, whereas Rab27B is observed in perinuclear vesicles. Targeted knockdown studies suggest that the isoforms control distinct steps of the exosome secretion pathway (6). Rab27A participates in docking and membrane fusion from multivesicular endosomes (MVEs) to exosomes, whereas Rab27B mediates the transfer of membrane vesicles from the trans-Golgi network to MVEs. It has been suggested that Rab27A and Rab27B may have importance in exosome secretion by cancer cells (7).				
Background References		1. Fukuda, M. (2013) <i>Traffic</i> 14, 949-63. 2. Izumi, T. (2007) <i>Endocr J</i> 54, 649-57. 3. Elstak, E.D. et al. (2011) <i>Blood</i> 118, 1570-8. 4. Bahadoran, P. et al. (2001) <i>J Cell Biol</i> 152, 843-50. 5. Wood, S.M. et al. (2009) <i>Blood</i> 114, 4117-27. 6. Ostrowski, M. et al. (2010) <i>Nat Cell Biol</i> 12, 19-30; sup pp 1-13. 7. Li, Z. et al. (2018) <i>Cell Commun Signal</i> 16, 44.				

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X

TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key W: Western Blotting

Cross-Reactivity Key H: Human

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for

more information.

Limited Uses Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST,

the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.