#4528 store at +4C

Pan-Keratin (C11) Mouse mAb (Alexa Fluor[®] 647 Conjugate)



Orders: 877-616-CELL (2355) orders@cellsignal.com Support: 877-678-TECH (8324)

Web:		

info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: IHC-P, IF-F, IF-IC, FC- FP	Reactivity: H M R Mk	Sensitivity: Endogenous	Source/Isotype: Mouse IgG1	UniProt ID: #P48668, #P13645, #P04259, #P05787, #P13646, #P02538, #P05783, #P13647, #P19013	Entrez-Gene Id: 286887, 3858, 3854, 3856, 3860, 3853, 3875, 3852, 3851	
Product Usage Information		Application Immunohistochemistry (Paraffin) Immunofluorescence (Frozen) Immunofluorescence (Immunocytochemistry) Flow Cytometry (Fixed/Permeabilized)		Dilution 1:50 - 1:200 1:100 - 1:400 1:800 - 1:3200 1:800		
Storage		Supplied in PBS (pH 7.2), less than 0.1% sodium azide and 2 mg/ml BSA. Store at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.				
Specificity/Sensit	ivity	Pan-Keratin (C11) Mouse mAb (Alexa Fluor [®] 647 Conjugate) detects endogenous levels of total keratins 4, 5, 6, 8, 10, 13 and 18. The antibody does not cross-react with other keratins.				
Source / Purificat	tion	Monoclonal antibody is produced by immunizing animals with a cytoskeleton preparation from A-431 cells. This antibody was conjugated to Alexa Fluor [®] 647 under optimal conditions with an F/P ratio of 2- 6. The Alexa Fluor [®] 647 dye is maximally excited by red light (e.g. 633 nm He-Ne laser). Antibody conjugates of the Alexa Fluor [®] 647 dye produce bright far-red-fluorescence emission, with a peak at 665 nm.				
Description		This Cell Signaling Technology [®] antibody is conjugated to Alexa Fluor [®] 647 fluorescent dye under optimal conditions. This antibody conjugate is expected to exhibit the same species cross-reactivity as the unconjugated Pan-Keratin (C11) Mouse mAb #4545.				
Background		Keratins (cytokeratins) are intermediate filament proteins that are mainly expressed in epithelial cells. Keratin heterodimers composed of an acidic keratin (or type I keratin, keratins K9-K28) and a basic keratin (or type II keratin, keratins K1-K8 and K71-K80) assemble to form filaments. Keratin isoforms demonstrate tissue- and differentiation-specific profiles that make them useful as research and clinical biomarkers (1,2).				
		Dysregulation/mutations in keratin genes can lead to a variety of disorders affecting the skin, hair, nails, and other epithelial tissues (3). While expression of keratins can be variable, immunohistochemical staining of keratins is widely used to help in the identification and classification of epithelial tumors, and may also provide prognostic information.				
		Keratins 8 and 18 (K8/K18) are expressed in simple epithelia of normal tissue, as well as in adenocarcinomas of the breast, lung, ovary, and gastrointestinal tract. Keratin 17 is expressed in basal keratinocytes of stratified epithelia, hair follicles, and sebaceous glands. Onset of keratin 17 expression coincides with the definition of major epithelial lineages during skin development (4). Keratin 14 (K14) is expressed in basal cells of stratified epithelia, and in basal-like subtypes of breast cancer and squamous cell carcinomas. Keratin 19 (K19) is expressed in glandular epithelia, including the liver, gallbladder, and pancreas, as well as in adenocarcinomas of the breast, thyroid, and bile duct. Keratin 20 (K20) is expressed in gastrointestinal epithelium, urothelium, and Merkel cells in the skin, as well as in colorectal carcinomas and some urothelial carcinomas. Keratin 5/6 (K5/6) is expressed in basal cells of stratified epithelia, including the skin, prostate, and breast, as well as in basal-like breast cancers, squamous cell carcinomas, and some lung carcinomas. Keratin 7 (K7) is expressed in glandular epithelia, such as those in the lung, breast, and female reproductive tract, as well as in adenocarcinomas of the lung, breast, and ovary (5,6).				
Keratins, particularly K8, K18, and K19, serve as biomarkers for identification of circul (CTCs) (5).					cation of circulating tumor cells	

Post-translational modifications, including phosphorylation, acetylation, ubiquitylation, sumoylation, glycosylation, and transamidation, have been shown to affect the functions of keratins in normal and disease states (6). Understanding the molecular mechanisms underlying these PTMs may provide insights into cancer pathogenesis.			
1. Chang, L. and Goldman, R.D. (2004) <i>Nat Rev Mol Cell Biol</i> 5, 601-13.			
2. Schweizer, J. et al. (2006) <i>J Cell Biol</i> 174, 169-74.			
3. Sarma, A. (2022) <i>Int J Biol Macromol</i> 219, 395-413. 4. McGowan, K.M. and Coulombe, P.A. (1998) <i>J Cell Biol</i> 143, 469-86.			
5. Werner, S. et al. (2020) <i>Mol Aspects Med</i> 72, 100817.			
6. Dmello, C. et al. (2019) <i>J Biosci</i> 44, 33.			
Species reactivity is determined by testing in at least one approved application (e.g., western blot).			
IHC-P: Immunohistochemistry (Paraffin) IF-F: Immunofluorescence (Frozen) IF-IC: Immunofluorescence (Immunocytochemistry) FC-FP: Flow Cytometry (Fixed/Permeabilized)			
H: Human M: Mouse R: Rat Mk: Monkey			
Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.			
This product is provided under an intellectual property license from Life Technologies Corporation. The transfer of this product is conditioned on the buyer using the purchased product solely in research conducted by the buyer, excluding contract research or any fee for service research, and the buyer must not (1) use this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; or (c) manufacturing or quality assurance or quality control, and/or (2) sell or transfer this product or its product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com.			
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.			
Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.			
Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with			