## 4629

## TBC1D1 (V796) Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:	
W, IP	M	Endogenous	160	Rabbit	#Q86TI0	23216	
Product Usage Information	2	<b>Application</b> Western Blotting Immunoprecipitation			<b>Dilution</b> 1:1000 1:50		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.					
Specificity/Sensitivity		TBC1D1 (V796) Antibody detects endogenous levels of total TBC1D1 protein.					
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to the sequence around Val796 of mouse TBC1D1. Antibodies are purified by protein A and peptide affinity chromatography.					
Background		TBC1D1 is a paralog of AS160 (1) and both proteins share about 50% identity (2). TBC1D1 was shown to be a candidate gene for severe obesity (3). It plays a role in Glut4 translocation through its GAP activity (2,4). Studies indicate that TBC1D1 is highly expressed in skeletal muscle (1). Insulin, AICAR, and contraction directly regulate TBC1D1 phosphorylation in this tissue (1). Three AMPK phosphorylation sites (Ser231, Ser660, and Ser700) and one Akt phosphorylation site (Thr590) were identified in skeletal muscle (5). Muscle contraction or AICAR treatment increases phosphorylation on Ser231, Ser660, and Ser700 but not on Thr590; insulin increases phosphorylation on Thr590 only (5).					
Background References		1. Taylor, E.B. et al. (2008) <i>J Biol Chem</i> 283, 9787-96. 2. Roach, W.G. et al. (2007) <i>Biochem J</i> 403, 353-8. 3. Stone, S. et al. (2006) <i>Hum Mol Genet</i> 15, 2709-20. 4. Chavez, J.A. et al. (2008) <i>J Biol Chem</i> 283, 9187-95. 5. Vichaiwong, K. et al. (2010) <i>Biochem J</i> 431, 311-20.					
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).					
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.					
Applications Key		W: Western Blotting IP: Immunoprecipitation					
Cross-Reactivity Key		M: Mouse					
<b>Trademarks and Patents</b>		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.					
		All other trademarks a more information.	are the property of t	heir respective owners.	Visit cellsignal.com	trademarks for	
Limited Uses		Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.					
			Products are labeled with For Research Use Only or a similar labeling statement and have not been				

approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a

component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com

For Research Use Only. Not for Use in Diagnostic Procedures.