

## YY1 (D5D9Z) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

info@cellsignal.com cellsignal.com Web:

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

10 <sup>6</sup> cells) per IP. This The CUT&RUN dilutio	antibody has been van was determined us was determined us was determined us a community of the community of	se 10 µl of antibody and validated using SimpleCh Ising CUT&RUN Assay Ki sing CUT&Tag Assay Kit #	nIP <sup>®</sup> Enzymatic Chr t #86652.	
The CUT&Tag dilution  Application  Western Blotting  Immunoprecipitation  Immunofluorescence Flow Cytometry (Fixe Chromatin IP	n was determined us n e (Immunocytochem	-		
Application Western Blotting Immunoprecipitation Immunofluorescence Flow Cytometry (Fixe Chromatin IP	n e (Immunocytochen	sing CUT&Tag Assay Kit #	‡77552.	
Western Blotting Immunoprecipitation Immunofluorescence Flow Cytometry (Fixe Chromatin IP	(Immunocytochem			
CUT&RUN CUT&Tag		istry)		Dilution 1:1000 1:200 1:800 1:100 1:50 1:50 1:50
		5), 150 mM NaCl, 100 μg/ ot aliquot the antibody.	/ml BSA, 50% glyce	rol and less than
YY1 (D5D9Z) Rabbit mAb recognizes endogenous levels of total YY1 protein.				
carboxy terminus of l	numan YY1 protein.	nunizing animals with re The antibody epitope ha Pro284 of human YY1 pro	as been further ma	
YY1 (Yin Yang1) is a ubiquitously expressed transcription factor with fundamental roles in embryogenesis, differentiation, replication and proliferation. YY1 contains four zinc finger motifs of the Cys-Cys-His-His type and can activate different eukaryotic genes (such as CREB, c-myc, Histone H4, p53 and PARP-1) or repress different eukaryotic genes (such as $\alpha$ -actin, IFN- $\beta$ and IFN- $\gamma$ ) as well as regulate some viral promoters (1). YY1 deficient embryos die approximately at the time of implantation, suggesting that YY1 has an essential role in embryonic development (2). YY1 is overexpressed in cancer cells such as prostate cancer and therefore may be considered a prognostic marker (1).				
1. Gordon, S. et al. (2006) <i>Oncogene</i> 25, 1125-42. 2. Donohoe, M.E. et al. (1999) <i>Mol. Cell Biol.</i> 19, 7237-7244.				
Species reactivity is d	etermined by testin	g in at least one approve	ed application (e.g.,	, western blot).
IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
W: Western Blotting IP: Immunoprecipitation IF-IC: Immunofluorescence (Immunocytochemistry) FC-FP: Flow Cytometry (Fixed/Permeabilized) ChIP: Chromatin IP ChIP-seq: Chromatin IP-seq C&R: CUT&RUN C&T: CUT&Tag				
<b>H:</b> Human <b>M:</b> Mouse	R: Rat Mk: Monkey			
Cell Signaling Techno	logy is a trademark	of Cell Signaling Techno	logy, Inc.	
	cells such as prostate  1. Gordon, S. et al. (20 2. Donohoe, M.E. et a  Species reactivity is d  IMPORTANT: For wes TBS, 0.1% Tween® 20  W: Western Blotting 1  FP: Flow Cytometry (F CUT&RUN C&T: CUT&  H: Human M: Mouse	cells such as prostate cancer and therefo  1. Gordon, S. et al. (2006) Oncogene 25, 1  2. Donohoe, M.E. et al. (1999) Mol. Cell Bi  Species reactivity is determined by testin  IMPORTANT: For western blots, incubate TBS, 0.1% Tween® 20 at 4°C with gentle s  W: Western Blotting IP: Immunoprecipita FP: Flow Cytometry (Fixed/Permeabilized CUT&RUN C&T: CUT&Tag  H: Human M: Mouse R: Rat Mk: Monkey	cells such as prostate cancer and therefore may be considered a  1. Gordon, S. et al. (2006) Oncogene 25, 1125-42.  2. Donohoe, M.E. et al. (1999) Mol. Cell Biol. 19, 7237-7244.  Species reactivity is determined by testing in at least one approve IMPORTANT: For western blots, incubate membrane with diluted TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.  W: Western Blotting IP: Immunoprecipitation IF-IC: Immunofluor FP: Flow Cytometry (Fixed/Permeabilized) ChIP: Chromatin IP Chi CUT&RUN C&T: CUT&Tag  H: Human M: Mouse R: Rat Mk: Monkey	cells such as prostate cancer and therefore may be considered a prognostic marker  1. Gordon, S. et al. (2006) <i>Oncogene</i> 25, 1125-42.  2. Donohoe, M.E. et al. (1999) <i>Mol. Cell Biol.</i> 19, 7237-7244.  Species reactivity is determined by testing in at least one approved application (e.g., IMPORTANT: For western blots, incubate membrane with diluted primary antibody i TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.  W: Western Blotting IP: Immunoprecipitation IF-IC: Immunofluorescence (Immunoc FP: Flow Cytometry (Fixed/Permeabilized) ChIP: Chromatin IP ChIP-seq: Chromatin CUT&RUN C&T: CUT&Tag

Alexa Fluor is a registered trademark of Life Technologies Corporation.

The transfer of this product is contingent on the buyer using the purchased product solely in research conducted by the buyer (whether the buyer is an academic or for-profit entity), for Immunocytochemistry, high content screening (HCS) analysis, or flow cytometry applications. The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) resale, whether or not such product or its components are resold for use in research; or for any other commercial purpose. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

## **Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.