TESK1 (D49D4) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 68	Source/Isotype: Rabbit IgG	UniProt ID: #Q15569	Entrez-Gene Id: 7016	
Product Usage Information		Application Western Blotting Immunoprecipitation			Dilution 1:1000 1:200		
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.					
Specificity/Sensitivity		TESK1 (D49D4) Rabbit mAb detects endogenous levels of total TESK1 protein.					
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to carboxy terminal residues of human TESK1 protein.					
Background		Testis-specific kinase 1 (TESK1) is a LIMK-related protein kinase originally identified to be expressed highly in testes and subsequently shown to be expressed in a wide variety of tissues and cell types (1-4). TESK1 phosphorylates the actin severing protein cofilin at Ser3, inactivating cofilin and thus regulating the organization of the actin cytoskeleton (2). Integrin signaling activates TESK1 activity and leads to stress fiber formation and cell spreading (2,5,6). TESK1 is involved in regulation of ERK signaling through its interaction with Spry2 (7) and regulation of cell spreading through its interaction with the focal adhesion protein actopaxin/α-parvin (5).					
Background References		 Toshima, J. et al. (1995) J Biol Chem 270, 31331-7. Toshima, J. et al. (2001) Mol Biol Cell 12, 1131-45. Toshima, J. et al. (2001) J Biol Chem 276, 31449-58. Toshima, J. et al. (2001) Biochem Biophys Res Commun 286, 566-73. LaLonde, D.P. et al. (2005) J Biol Chem 280, 21680-8. Tsumura, Y. et al. (2005) Biochem J 387, 627-37. Chandramouli, S. et al. (2008) J Biol Chem 283, 1679-91. 					

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key W: Western Blotting IP: Immunoprecipitation

Cross-Reactivity Key H: Human

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. Trademarks and Patents

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for

more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for

diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.