

WRN (8H3) Mouse mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, W-S, IF-IC	H M	Endogenous	200	Mouse IgG1	#Q14191	7486

Product Usage Information**Application**

Western Blotting
Simple Western™
Immunofluorescence (Immunocytochemistry)

Dilution

1:1000
1:10 - 1:50
1:1600 - 1:6400

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

WRN (8H3) Mouse mAb detects endogenous levels of total WRN protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with full length human recombinant WRN protein.

Background

The RecQ family of DNA and RNA helicases is a family of enzymes that has been shown to be important to genome integrity (1). Members of this family function in several DNA repair processes, including double-strand break repair, homologous recombination, and re-initiation of DNA replication at stalled replication forks (2,3). Mutations in RecQ helicase family members result in syndromes that display varying types of chromosomal abnormalities and overall genomic instability (1). *WRN* is a member of the RecQ family that has been identified as the gene underlying Werner's syndrome, an autosomal recessive disorder characterized by premature aging and predisposition to cancer (4). Cells from Werner's syndrome patients exhibit genomic instability that is associated with deficient DNA repair activity (5).

Background References

1. Cobb, J.A. et al. (2002) *FEBS Lett* 529, 43-8.
2. Ouyang, K.J. et al. *Mech Ageing Dev* 129, 425-40.
3. Kaneko, H. et al. (2004) *Adv Biophys* 38, 45-64.
4. Epstein, C.J. et al. (1966) *Medicine (Baltimore)* 45, 177-221.
5. Salk, D. et al. (1985) *Adv Exp Med Biol* 190, 541-6.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **W-S:** Simple Western™ **IF-IC:** Immunofluorescence (Immunocytochemistry)

Cross-Reactivity Key

H: Human **M:** Mouse

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in

any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.