

47134

XBP-1s (E7M5C) Mouse mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IF-IC, FC-FP	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 60	Source/Isotype: Mouse IgG1	UniProt ID: #P17861-2	Entrez-Gene Id: 7494
Product Usage Information		Application Western Blotting Immunofluorescence Flow Cytometry (Fixed	•	istry)		Dilution 1:1000 1:100 1:200
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody. For a carrier free (BSA and azide free) version of this product see product #30163.				
Specificity/Sensitivity		XBP-1s (E7M5C) Mouse mAb recognizes endogenous levels of total XBP-1s protein.				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Val224 of human XBP-1s protein.				
Background	Following protein synthesis, secretory, intra-organellar, and transmembrane proteins translocate into the endoplasmic reticulum (ER) where they are post-translationally modified and properly folded. The accumulation of unfolded proteins within the ER triggers an adaptive mechanism known as the unfolded protein response (UPR) that counteracts compromised protein folding (1). The transmembrane serine/threonine kinase IRE1, originally identified in <i>Saccharomyces cerevisiae</i> , is a proximal sensor for the UPR that transmits the unfolded protein signal across the ER membrane (2-4). The human homolog IRE1a was later identified and is ubiquitously expressed in human tissues (5). Upon activation of the unfolded protein response, IRE1a splices X-box binding protein 1 (XBP-1) mRNA through an unconventional mechanism using its endoribonuclease activity (6). This reaction converts XBP-1 from an unspliced XBP-1u isoform to the spliced XBP-1s isoform, which is a potent transcriptional activator that induces expression of many UPR responsive genes (6).					
1. Kaufman, R.J. et al. (2002) Nat Rev Mol Cell Biol 3, 411-21. 2. Nikawa, J. and Yamashita, S. (1992) Mol Microbiol 6, 1441-6. 3. Cox, J.S. et al. (1993) Cell 73, 1197-206. 4. Mori, K. et al. (1993) Cell 74, 743-56. 5. Tirasophon, W. et al. (1998) Genes Dev 12, 1812-24. 6. Lee, K. et al. (2002) Genes Dev 16, 452-66.						
Species Reactiv	/ity	Species reactivity is d	etermined by testin	g in at least one approve	ed application (e.g.,	western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X

TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting IF-IC: Immunofluorescence (Immunocytochemistry) FC-FP: Flow Cytometry

(Fixed/Permeabilized)

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Alexa Fluor is a registered trademark of Life Technologies Corporation.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.