

#4749 Store at -20°C

# Integrin Antibody Sampler Kit



✓ 1 Kit  
(7 x 20 µl)

**Orders** ■ 877-616-CELL (2355)  
orders@cellsignal.com  
**Support** ■ 877-678-TECH (8324)  
info@cellsignal.com  
**Web** ■ www.cellsignal.com

rev. 06/16

**For Research Use Only. Not For Use In Diagnostic Procedures.**

Products Included	Product #	Quantity	Mol. Wt.	Source
Integrin α4 (D2E1) XP® Rabbit mAb	8440	20 µl	70, 140, 150 kDa	Rabbit IgG
Integrin α5 Antibody	4705	20 µl	150 kDa	Rabbit IgG
Integrin αV Antibody	4711	20 µl	135, 140 kDa	Rabbit IgG
Integrin β1 (D2E5) Rabbit mAb	9699	20 µl	115, 135 kDa	Rabbit IgG
Integrin β3 (D7X3P) XP® Rabbit mAb	13166	20 µl	100 kDa	Rabbit IgG
Integrin β4 (D8P6C) XP® Rabbit mAb	14803	20 µl	210 kDa	Rabbit IgG
Integrin β-5 (D24A5) Rabbit mAb	3629	20 µl	90 kDa	Rabbit IgG
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat

**Storage:** Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. *Do not aliquot the antibodies.*

**Recommended Antibody Dilutions:**  
Western blotting 1:1000

**Please visit [www.cellsignal.com](http://www.cellsignal.com) for validation data and a complete listing of recommended companion products.**

See [www.cellsignal.com](http://www.cellsignal.com) for individual component applications, species cross-reactivity, dilutions and additional application protocols.

**Description:** The Integrin Antibody Sampler Kit provides an economical means to screen samples for α and β subunits of integrin molecules. The kit includes enough antibody to perform two western blot experiments with each primary antibody.

**Background:** Integrins are α/β heterodimeric cell surface receptors that play a pivotal role in cell adhesion and migration, as well as in growth and survival (1,2). The integrin family contains at least 18 α and 8 β subunits that form 24 known integrins with distinct tissue distribution and overlapping ligand specificities (3). Integrins not only transmit signals to cells in response to the extracellular environment (outside-in signaling), but also sense intracellular cues to alter their interaction with the extracellular environment (inside-out signaling) (1,2).

**Specificity/Sensitivity:** All antibodies in the Integrin Antibody Sampler Kit detect endogenous levels of their target α and β integrin subunits of the integrin dimer. None of the antibodies provided detect intact dimers or their activation state.

**Source/Purification:** Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ser458 of human integrin α5 protein, Arg108 of human integrin αV. Antibodies are purified by protein A and peptide affinity chromatography. Monoclonal antibodies are produced by immunizing animals with a synthetic peptide sequence corresponding to residues surrounding Ser1027 of human integrin α4 protein, the carboxy terminus of human integrin β4 protein, Pro680 of human integrin β1 protein, Ile114 of human integrin β3 protein, and Lys789 of human β5 protein.

**Background References:**  
(1) Liu, S. et al. (2000) *J Cell Sci* 113 ( Pt 20), 3563–71.  
(2) Hood, J.D. and Cheresch, D.A. (2002) *Nat Rev Cancer* 2, 91–100.  
(3) van der Flier, A. and Sonnenberg, A. (2001) *Cell Tissue Res* 305, 285–98.

© 2016 Cell Signaling Technology, Inc. XP® and Cell Signaling Technology® are trademarks of Cell Signaling Technology, Inc.

**Applications Key:** W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide  
**Species Cross-Reactivity Key:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine  
Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.

## Western Immunoblotting Protocol

For western blots, incubate membrane with diluted primary antibody in either 5% w/v BSA or nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**NOTE:** Please refer to primary antibody datasheet or product webpage for recommended primary antibody dilution buffer and recommended antibody dilution.

### A. Solutions and Reagents

**NOTE:** Prepare solutions with reverse osmosis deionized (RODI) or equivalent grade water.

- 20X Phosphate Buffered Saline (PBS):** (#9808) To prepare 1 L 1X PBS: add 50 ml 20X PBS to 950 ml dH<sub>2</sub>O, mix.
- 10X Tris Buffered Saline (TBS):** (#12498) To prepare 1 L 1X TBS: add 100 ml 10X to 900 ml dH<sub>2</sub>O, mix.
- 1X SDS Sample Buffer:** Blue Loading Pack (#7722) or Red Loading Pack (#7723)  
Prepare fresh 3X reducing loading buffer by adding 1/10 volume 30X DTT to 1 volume of 3X SDS loading buffer. Dilute to 1X with dH<sub>2</sub>O.
- 10X Tris-Glycine SDS Running Buffer:** (#4050) To prepare 1 L 1X running buffer: add 100 ml 10X running buffer to 900 ml dH<sub>2</sub>O, mix.
- 10X Tris-Glycine Transfer Buffer:** (#12539) To prepare 1 L 1X transfer buffer: add 100 ml 10X transfer buffer to 200 ml methanol + 700 ml dH<sub>2</sub>O, mix.
- 10X Tris Buffered Saline with Tween® 20 (TBST):** (#9997) To prepare 1 L 1X TBST: add 100 ml 10X TBST to 900 ml dH<sub>2</sub>O, mix.
- Nonfat Dry Milk:** (#9999)
- Blocking Buffer:** 1X TBST with 5% w/v nonfat dry milk; for 150 ml, add 7.5 g nonfat dry milk to 150 ml 1X TBST and mix well.
- Wash Buffer:** (#9997) 1X TBST
- Bovine Serum Albumin (BSA):** (#9998)
- Primary Antibody Dilution Buffer:** 1X TBST with 5% BSA or 5% nonfat dry milk as indicated on primary antibody datasheet; for 20 ml, add 1.0 g BSA or nonfat dry milk to 20 ml 1X TBST and mix well.
- Biotinylated Protein Ladder Detection Pack:** (#7727)
- Prestained Protein Marker, Broad Range (Premixed Format):** (#7720)
- Blotting Membrane and Paper:** (#12369) This protocol has been optimized for nitrocellulose membranes. Pore size 0.2 µm is generally recommended.
- Secondary Antibody Conjugated to HRP:** anti-rabbit (#7074); anti-mouse (#7076)
- Detection Reagent:** LumiGLO® chemiluminescent reagent and peroxide (#7003) or SignalFire™ ECL Reagent (#6883)

### B. Protein Blotting

**A general protocol for sample preparation.**

- Treat cells by adding fresh media containing regulator for desired time.
- Aspirate media from cultures; wash cells with 1X PBS; aspirate.
- Lyse cells by adding 1X SDS sample buffer (100 µl per well of 6-well plate or 500 µl for a 10 cm diameter plate). Immediately scrape the cells off the plate and transfer the extract to a microcentrifuge tube. Keep on ice.
- Sonicate for 10–15 sec to complete cell lysis and shear DNA (to reduce sample viscosity).
- Heat a 20 µl sample to 95–100°C for 5 min; cool on ice.
- Microcentrifuge for 5 min.
- Load 20 µl onto SDS-PAGE gel (10 cm x 10 cm). **NOTE:** Loading of prestained molecular weight markers (#7720, 10 µl/lane) to verify electrotransfer and biotinylated protein ladder (#7727, 10 µl/lane) to determine molecular weights are recommended.
- Electrotransfer to nitrocellulose membrane (#12369).

### C. Membrane Blocking and Antibody Incubations

**NOTE:** Volumes are for 10 cm x 10 cm (100 cm<sup>2</sup>) of membrane; for different sized membranes, adjust volumes accordingly.

#### I. Membrane Blocking

- (Optional) After transfer, wash nitrocellulose membrane with 25 ml TBS for 5 min at room temperature.
- Incubate membrane in 25 ml of blocking buffer for 1 hr at room temperature.
- Wash three times for 5 min each with 15 ml of TBST.

#### II. Primary Antibody Incubation

- Incubate membrane and primary antibody (at the appropriate dilution and diluent as recommended in the product datasheet) in 10 ml primary antibody dilution buffer with gentle agitation overnight at 4°C.
- Wash three times for 5 min each with 15 ml of TBST.
- Incubate membrane with the species appropriate HRP-conjugated secondary antibody (#7074 or #7076 at 1:2000) and anti-biotin, HRP-linked Antibody (#7075 at 1:1000–1:3000) to detect biotinylated protein markers in 10 ml of blocking buffer with gentle agitation for 1 hr at room temperature.
- Wash three times for 5 min each with 15 ml of TBST.
- Proceed with detection (Section D).

### D. Detection of Proteins

- Incubate membrane with 10 ml LumiGLO® (0.5 ml 20X LumiGLO® #7003, 0.5 ml 20X peroxide, and 9.0 ml purified water) or 10 ml SignalFire™ #6883 (5 ml Reagent A, 5 ml Reagent B) with gentle agitation for 1 min at room temperature.
- Drain membrane of excess developing solution (do not let dry), wrap in plastic wrap and expose to x-ray film. An initial 10 sec exposure should indicate the proper exposure time.  
**NOTE:** Due to the kinetics of the detection reaction, signal is most intense immediately following incubation and declines over the following 2 hr.

LumiGLO® is a registered trademark of Kirkegaard & Perry Laboratories. Tween® is a registered trademark of ICI Americas, INC. SignalFire™ is a trademark of Cell Signaling Technology, INC.

Orders ■ 877-616-CELL (2355) orders@cellsignal.com Support ■ 877-678-TECH (8324) info@cellsignal.com Web ■ www.cellsignal.com