DCBLD2 Antibody



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Applications: W	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 120	Source/Isotype: Rabbit	UniProt ID: #Q96PD2	Entrez-Gene Id: 131566
Product Usage Information		Application Western Blotting			Dilution 1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA and 50% glycerol. Store at – 20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		DCBLD2 Antibody detects endogenous levels of total DCBLD2 protein.				
Species predicted to react based on 100% sequence homology		Monkey				
Source / Purification		Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human DCBLD2 protein. Antibodies are purified by protein A and peptide affinity chromatography.				
Background		DCBLD2 (discoidin, CUB and LCCL domain-containing 2; also known as ESDN and CLCP1) is a type I transmembrane protein that is structurally similar to neuropilin family proteins and contains the longest known amino-terminal secretory signal sequence among eukaryotic proteins (1). Highly expressed in nerve bundles, vascular smooth muscle cells and upregulated following vascular injury, DCBLD2 may be involved in a wide range of functions in the nervous and vascular systems (1,2). Studies have found DCBLD2 to be upregulated in several types of lung cancer with an especially high frequency in metastatic lesions and lymph node metastasis, implicating a role in the process of tumor progression and metastatic capability (3). DCBLD2 has also been identified as part of a complex EGF phosphotyrosine signaling network, serving as a novel tyrosine phosphorylation target of EGF signaling in human cancer cells (4).				
Background References		1. Kobuke, K. et al. (2001) <i>J Biol Chem</i> 276, 34105-14. 2. Sadeghi, M.M. et al. (2007) <i>Am J Transplant</i> 7, 2098-105. 3. Koshikawa, K. et al. (2002) <i>Oncogene</i> 21, 2822-8. 4. Chen, Y. et al. (2007) <i>Proteomics</i> 7, 2384-97.				
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).				
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				
Applications Key		W: Western Blotting				
Cross-Reactivity Key		H: Human M: Mouse R: Rat				
Trademarks and Patents		Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.				
		All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.				

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.