



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

#48127 store at +4C

IL2-R α /CD25 (PC61.5) Rat mAb (violetFluor™ 450 Conjugate)

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
FC-L	M	Endogenous	Rat IgG1 lambda	#P01590	16184
Product Usage Information	For optimal flow cytometry results, we recommend 0.5 μ g of antibody per test.				
	Application				Dilution
	Flow Cytometry (Live)				1:40
Storage	Supplied in 10 mM NaH ₂ PO ₄ , 150 mM NaCl, 0.09% NaN ₃ , 0.1% gelatin, pH 7.2. This product is stable for 12 months when stored at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze.				
Specificity/Sensitivity	IL2-R α /CD25 (PC61.5) Rat mAb (violetFluor™ 450 Conjugate) recognizes endogenous levels of total IL2-R α /CD25 protein. This antibody detects an epitope within the extracellular domain.				
Source / Purification	This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation.				
Description	This Cell Signaling Technology antibody is conjugated to violetFluor™ 450 and tested in-house for direct flow cytometric analysis in mouse cells.				
Background	Interleukin-2 (IL-2) is a T cell stimulatory cytokine best known for inducing T cell proliferation and NK cell proliferation and activation (1,2). IL-2 also promotes peripheral development of regulatory T cells (Tregs) (3,4). Conversely, IL-2 is involved in the activation-induced cell death (AICD) that is observed post T cell expansion by increasing levels of Fas on CD4 ⁺ T cells (5). The effects of IL-2 are mediated through a trimeric receptor complex consisting of IL-2R α , IL-2R β , and the common gamma chain, γ c (1,2). IL-2R α binds exclusively to IL-2 with low affinity and increases the binding affinity of the whole receptor complex including IL-2R β and γ c subunits. IL-15 also binds to IL-2R β (1,2). γ c is used by other cytokines, including IL-4, IL-7, IL-9, IL-15, and IL-21 (1,2). Binding of IL-2 initiates signaling cascades involving Jak1, Jak3, Stat5, and the PI3K/Akt pathways (1,2).				
Background References	<ol style="list-style-type: none"> Ma, A. et al. (2006) <i>Annu Rev Immunol</i> 24, 657-79. Gaffen, S.L. and Liu, K.D. (2004) <i>Cytokine</i> 28, 109-23. Fehérvári, Z. et al. (2006) <i>Trends Immunol</i> 27, 109-11. Antony, P.A. et al. (2006) <i>J Immunol</i> 176, 5255-66. Jaleco, S. et al. (2003) <i>J Immunol</i> 171, 61-8. 				

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key **FC-L:** Flow Cytometry (Live)

Cross-Reactivity Key **M:** Mouse

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in

any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.