

Store at
-80°C

#48801

SARS-CoV-2 Spike RBD (318-541) Recombinant Protein (8xHis-Tag)

100 µg



Support: +1-978-867-2388 (U.S.)
www.cellsignal.com/support

Orders: 877-616-2355 (U.S.)
orders@cellsignal.com


Entrez-Gene ID #43740568
UniProt ID #P0DTC2

rev. 10/19/20

For Research Use Only. Not For Use In Diagnostic Procedures.

Description: SARS-CoV-2 Spike RBD (318-541) Recombinant Protein (8xHis-Tag) is derived from a recombinant expression construct corresponding to the host receptor-binding domain (RBD) of the SARS-CoV-2 spike protein. The expressed protein contains an 8xHis-Tag at its carboxy terminus.

Background: The cause of the COVID-19 pandemic is a novel and highly pathogenic coronavirus, termed SARS-CoV-2 (severe acute respiratory syndrome coronavirus-2). SARS-CoV-2 is a member of the Coronaviridae family of viruses (1). The genome of SARS-CoV-2 is similar to other coronaviruses, and is comprised of four key structural proteins: S, the spike protein, E, the envelope protein, M, the membrane protein, and N, the nucleocapsid protein (2). Coronavirus spike proteins are class I fusion proteins and harbor an ectodomain, a transmembrane domain, and an intracellular tail (3,4). The highly glycosylated ectodomain projects from the viral envelope surface and facilitates attachment and fusion with the host cell plasma membrane. The ectodomain can be further subdivided into host receptor-binding domain (RBD) (S1) and membrane-fusion (S2) subunits, which are produced upon proteolysis by host proteases at S1/S2 and S2' sites. S1 and S2 subunits remain associated after cleavage and assemble into crown-like homotrimers (2,4). In humans, both SARS-CoV and SARS-CoV-2 spike proteins utilize the angiotensin-converting enzyme 2 (ACE2) protein as a receptor



The purity of SARS-CoV-2 Spike RBD (318-541) Recombinant Protein (8xHis-Tag) was determined by densitometry after SDS-PAGE of 2 µg of protein, followed by staining with Coomassie Blue. Purity values were determined from the DTT-reduced samples (+).

for cellular entry (5-7). Spike protein subunits represent a key antigenic feature of coronavirus virions, and therefore represent an important target of vaccines, novel therapeutic antibodies, and small-molecule inhibitors (8,9).

Molecular Weight: 34, 68 kDa (non-reduced); 36 kDa (reduced)

Formulation:

Expression Host: Human (HEK293 cells)
Supplied in PBS solution (pH 7.2).

Purity: 98%, determined by SDS-PAGE.


Storage: Stable at -80°C for 1 year after receipt. Avoid repeated freeze-thaw cycles.

Background References:

- (1) Zhou, P. et al. (2020) *Nature* 579, 270-3.
- (2) Tortorici, M.A. and Veasler, D. (2019) *Adv Virus Res* 105, 93-116.
- (3) Li, F. et al. (2006) *J Virol* 80, 6794-800.
- (4) Li, F. (2016) *Annu Rev Virol* 3, 237-61.
- (5) Shang, J. et al. (2020) *Nature* 581, 221-4.
- (6) Wrapp, D. et al. (2020) *Science* 367, 1260-3.
- (7) Yan, R. et al. (2020) *Science* 367, 1444-8.
- (8) Yuan, Y. et al. (2017) *Nat Commun* 8, 15092.
- (9) Amanat, F. and Krammer, F. (2020) *Immunity* 52, 583-9.

Immobilized **Human ACE2 (18-652) Recombinant Protein (mFc-Tag) #83986**


In Solution **SARS-CoV-2 Spike RBD (318-541) Recombinant Protein (8xHis-Tag) – by Conc. (nM)**



Binding kinetics between Human ACE2 (18-652) Recombinant Protein (mFc-Tag) #83986 (immobilized) and SARS-CoV-2 Spike RBD (318-541) Recombinant Protein (8xHis-Tag) (in solution, at the indicated concentrations). The vertical red line (180 sec) indicates addition of PBS to induce dissociation. Binding was detected with an anti-mouse Fc biosensor. Values on y-axis indicate binding response signals recorded for 5 different concentrations of SARS-CoV-2 Spike RBD (318-541) Recombinant Protein (8xHis-Tag) (62.5, 125, 250, 500, and 1000 nM).

Immobilized **SARS-CoV-2 Spike Protein (CR3022) Mouse IgG2a mAb #24387**

In Solution **SARS-CoV-2 Spike RBD (318-541) Recombinant Protein (8xHis-Tag) – by Conc. (nM)**



Binding kinetics between SARS-CoV-2 Spike Protein (CR3022) Mouse IgG2a mAb #24387 (immobilized) and SARS-CoV-2 Spike RBD (318-541) Recombinant Protein (8xHis-Tag) (in solution, at indicated concentrations). The vertical red line (180 sec) indicates addition of PBS to induce dissociation. Binding was detected with an anti-mouse Fc biosensor. Values on y-axis indicate binding response signals recorded for 5 different concentrations of SARS-CoV-2 Spike RBD (318-541) Recombinant Protein (8xHis-Tag) (31.2, 62.5, 125, 250, and 500 nM).

Thank you for your recent purchase. If you would like to provide a review visit cellsignal.com/comments.

www.cellsignal.com

© 2020 Cell Signaling Technology, Inc.

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Applications: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide Species Cross-Reactivity: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.