

Store at
-20°C

Ghost Dye Violet 450 Fixable Viability Dye



**Cell Signaling**
TECHNOLOGY®

#49826

100 µL

Support: +1-978-867-2388 (U.S.)
cellsignal.com/support**Orders:** 877-616-2355 (U.S.)
orders@cellsignal.com**For Research Use Only. Not for Use in Diagnostic Procedures.****Applications**
F**Species Cross-Reactivity**
All

Description: Ghost Dye Violet 450 Fixable Viability Dye is used to discriminate viable from non-viable mammalian cells in flow cytometry applications. Ghost Dye Violet 450 Fixable Viability Dye irreversibly binds free amines available on the cell surface as well as intracellular free amines exposed in cells with compromised cell membranes. Non-viable cells with loss of membrane integrity will be labeled with significantly more dye than healthy cells in the same sample. The cells may then be fixed, and the degree of labeling will be preserved through fixation, permeabilization, and antibody incubation steps. Cells that exhibit increased fluorescence intensity were non-viable at the time of fixation and can be excluded from analysis.

**Ghost Dye Violet 450****Ghost Dye Violet 450**

Flow cytometric analysis of live and fixed/permeabilized human peripheral blood mononuclear cells, combined and stained with Ghost Dye Violet 450 Fixable Viability Dye. Viable gate is indicated.

Storage: Store at -20°C desiccated and protected from light. This product is stable for 12 months. Aliquot to avoid excessive freeze-thaw cycles.

Directions For Use:

- Prepare the following reagents with reverse osmosis deionized (RODI) or equivalent grade water:
 - 1X PBS (azide- and protein/serum-free)
 - Incubation Buffer: Dissolve 0.5 g Bovine Serum Albumin (BSA) (#9998) in 100 ml 1X PBS. Store at 4°C.
- Remove Ghost Dye from -20°C and bring to room temperature.
- Collect cells by centrifugation and aspirate supernatant.
- Wash cells by centrifugation in excess 1X PBS. Repeat if necessary.
- Resuspend cells to a concentration of 1-10 x 10⁶/mL in 1X PBS.
- Centrifuge the Ghost Dye before opening then add 1 µL for each 1 mL of cell suspension and vortex immediately.
- Incubate for 30 minutes at 4°C protected from light.
- Wash by centrifugation in excess incubation buffer. Discard supernatant. Repeat.
- Cells can then be fixed, permeabilized and immunostained based upon experimental design and recommended protocols.
- Exclude cells with high Ghost Dye fluorescence from analysis. These were non-viable cells at the time of fixation. See details below for excitation and emission specifications.

Ghost Dye Violet 450 Fixable Viability Dye is excited by the violet (405 nm) laser line and has a peak emission of 450 nm that can be detected using 440/40 or 440/50 band pass filters commonly used for detection of violetFluor 450, Pacific Blue, etc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

cellsignal.com

© 2024 Cell Signaling Technology, Inc.

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Applications: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry CHIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry FC-FP—Flow cytometry-Fixed/Permeabilized FC-L—Flow cytometry-Live E-P—ELISA-Peptide
Species Cross-Reactivity: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse
All—all species expected. Species enclosed in parentheses are predicted to react based on 100% homology.