
Store at

50996

#





Orders:	877-616-CELL (2355) orders@cellsignal.com
Support:	877-678-TECH (8324)
Web:	info@cellsignal.com cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, W-S, IP	<b>Reactivity:</b> H M R	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 130	Source/Isotype: Mouse IgG2a	<b>UniProt ID:</b> #P23458	Entrez-Gene Id: 3716		
Product Usage Information		<b>Application</b> Western Blotting Simple Western™ Immunoprecipitation			<b>Dilution</b> 1:1000 1:50 - 1:250 1:200			
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.						
Specificity/Sen	sitivity	<b>ivity</b> Jak1 (D1T6W) Mouse mAb recognizes endogenous levels of total Jak1 protein.						
Source / Purifi	cation	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding His869 of human Jak1 protein.						
Background		Members of the Janus family of tyrosine kinases (Jak1, Jak2, Jak3, and Tyk2) are activated by ligands binding to a number of associated cytokine receptors (1). Upon cytokine receptor activation, Jak proteins become autophosphorylated and phosphorylate their associated receptors to provide multiple binding sites for signaling proteins. These associated signaling proteins, such as Stats (2), Shc (3), insulin receptor substrates (4), and focal adhesion kinase (FAK) (5), typically contain SH2 or other phosphotyrosine-binding domains. Activation of Jak kinases upon cytokine receptor binding is associated with tyrosine phosphorylation within their activation loops, including Tyr1034/1035 of Jak1, Tyr1007/1008 of Jak2, Tyr980/981 of Jak3, and Tyr1054/1055 of Tyk2. Many studies have indicated that various cytokine receptors have clear preferences that utilize distinct Jak family members. Aberrant regulation of Jak signaling is associated with a number of diseases, including myeloproliferative neoplasms, leukemia, and inflammatory disease (6).						
Background Ro	eferences	1. Leonard, W.J. and O'Shea, J.J. (1998) <i>Annu Rev Immunol</i> 16, 293-322. 2. Darnell, J.E. (1997) <i>Science</i> 277, 1630-5. 3. VanderKuur, J. et al. (1995) <i>J Biol Chem</i> 270, 7587-93. 4. Argetsinger, L.S. et al. (1995) <i>J Biol Chem</i> 270, 14685-92. 5. Zhu, T. et al. (1998) <i>J Biol Chem</i> 273, 10682-9. 6. Babon, J.J. et al. (2014) <i>Biochem J</i> 462, 1-13.						
Species Reacti	vity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).						
Western Blot E	Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.						
Applications K	ey	<b>W:</b> Western Blotting <b>W-S:</b> Simple Western <sup>™</sup> <b>IP:</b> Immunoprecipitation						
Cross-Reactivi	ty Key	H: Human M: Mouse R: Rat						
Trademarks ar	nd Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.						
		All other trademarks a more information.	demarks are the property of their respective owners. Visit cellsignal.com/trademarks for ation.					
Limited Uses		the following terms ap terms and conditions t	as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, owing terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's nd conditions that are in addition to, or different from, those contained herein, unless ely accepted in writing by a legally authorized representative of CST, are rejected and are of no • effect.					

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.