

## FAM129B Antibody



**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP, IF-IC, FC-FP	H Mk	Endogenous	95	Rabbit	#Q96TA1	64855

### Product Usage Information

#### Application

Western Blotting  
Immunoprecipitation  
Immunofluorescence (Immunocytochemistry)  
Flow Cytometry (Fixed/Permeabilized)

#### Dilution

1:1000  
1:50  
1:50  
1:50

### Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

### Specificity/Sensitivity

FAM129B Antibody recognizes endogenous levels of total FAM129B protein.

### Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Glu705 of human Fam129B protein. Antibodies were purified by protein A and peptide affinity chromatography.

### Background

FAM129B/Niban-like protein 1 (family with sequence similarity 129, member B) belongs to a poorly characterized family of Niban proteins that also includes FAM129A/Niban and FAM129C/Niban-like protein 2. FAM129A/Niban is implicated in the ER stress response and is upregulated at the protein level in thyroid carcinoma (1,2). FAM129C/Niban-like protein 2 is preferentially expressed in B-cells and is one of five biomarkers upregulated in whole blood from patients with colorectal carcinoma (3,4). FAM129B is broadly expressed and has been shown to be a downstream target of B-Raf in melanoma cells (5). Though FAM129B does not appear to regulate cell growth and division, phosphorylation of FAM129B by B-Raf is essential for the invasive potential of melanoma and non-melanoma cell lines (5). Deletion of FAM129B in melanoma cells significantly impairs B-Raf/MEK/Erk-dependent invasion into the extracellular matrix (5).

### Background References

1. Sun, G.D. et al. (2007) *Biochem Biophys Res Commun* 360, 181-7.
2. Matsumoto, F. et al. (2006) *Hum Pathol* 37, 1592-600.
3. Boyd, R.S. et al. (2003) *Leukemia* 17, 1605-12.
4. Han, M. et al. (2008) *Clin Cancer Res* 14, 455-60.
5. Old, W.M. et al. (2009) *Mol Cell* 34, 115-31.

### Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

### Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

### Applications Key

**W:** Western Blotting **IP:** Immunoprecipitation **IF-IC:** Immunofluorescence (Immunocytochemistry) **FC-FP:** Flow Cytometry (Fixed/Permeabilized)

### Cross-Reactivity Key

**H:** Human **Mk:** Monkey

### Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

### Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.