

TBK1/NAK (E9H5S) Mouse mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 84	Source/Isotype: Mouse IgG1	UniProt ID: #Q9UHD2	Entrez-Gene Id: 29110
	Application			Dilution	
Information	Western Blotting			1:1000	
	Fluorescent Western			1:1000	
	Immunoprecipitation			1:100	
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
	For a carrier free (BSA a	nd azide free) ver	sion of this product see	product #56401.	
sitivity	TBK1/NAK (E9H5S) Mouse mAb recognizes endogenous levels of total TBK1/NAK protein.				
ted to react sequence	Monkey				
cation	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Glu540 of human TBK1/NAK protein.				
	TBK1 (TANK-binding kinase 1)/NAK (NF-κB activating kinase) is an IκB kinase (IKK)-activating kinase and can activate IKK through direct phosphorylation (1). TBK1 was identified through association with the TRAF binding protein, TANK, and found to function upstream of NIK and IKK in the activation of NF-κB (2). TBK1 induces IκB degradation and NF-κB activity through IKKβ. TBK1 may mediate IKK and NF-κB activation in response to growth factors that stimulate PKCε activity (1). TBK1 plays a pivotal role in the activation of IRF3 in the innate immune response (3).				
eferences	1. Tojima, Y. et al. (2000) <i>Nature</i> 404, 778-82. 2. Pomerantz, J.L. and Baltimore, D. (1999) <i>EMBO J</i> 18, 6694-704. 3. Fitzgerald, K.A. et al. (2003) <i>Nat Immunol</i> 4, 491-6.				
	HMR sitivity ted to react sequence cation	Application Western Blotting Fluorescent Western Immunoprecipitation Supplied in 10 mM sodi 0.02% sodium azide. Sto For a carrier free (BSA a TBK1/NAK (E9H5S) Mou ted to react sequence Monoclonal antibody is residues surrounding G TBK1 (TANK-binding kin can activate IKK throug TRAF binding protein, T (2). TBK1 induces IKB de activation in response t activation of IRF3 in the eferences 1. Tojima, Y. et al. (2000 2. Pomerantz, J.L. and B	Application Western Blotting Fluorescent Western Immunoprecipitation Supplied in 10 mM sodium HEPES (pH 7.5 0.02% sodium azide. Store at -20°C. Do not for a carrier free (BSA and azide free) versitivity TBK1/NAK (E9H5S) Mouse mAb recognize Monkey Monoclonal antibody is produced by immaresidues surrounding Glu540 of human TTBK1 (TANK-binding kinase 1)/NAK (NF-KE can activate IKK through direct phosphory TRAF binding protein, TANK, and found to (2). TBK1 induces IKB degradation and NF activation in response to growth factors to activation of IRF3 in the innate immune references 1. Tojima, Y. et al. (2000) Nature 404, 778-2. Pomerantz, J.L. and Baltimore, D. (1999)	Application Western Blotting Fluorescent Western Immunoprecipitation Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody. For a carrier free (BSA and azide free) version of this product see TBK1/NAK (E9H5S) Mouse mAb recognizes endogenous levels of the sequence Monkey Monoclonal antibody is produced by immunizing animals with a sequence Monkey Monoclonal antibody is produced by immunizing animals with a sequence TBK1 (TANK-binding kinase 1)/NAK (NF-kB activating kinase) is an can activate IKK through direct phosphorylation (1). TBK1 was ide TRAF binding protein, TANK, and found to function upstream of N (2). TBK1 induces IkB degradation and NF-kB activity through IKK activation in response to growth factors that stimulate PKCe activactivation of IRF3 in the innate immune response (3). 1. Tojima, Y. et al. (2000) Nature 404, 778-82. 2. Pomerantz, J.L. and Baltimore, D. (1999) EMBO J 18, 6694-704.	Application Western Blotting Fluorescent Western Inmunoprecipitation Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glyce 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody. For a carrier free (BSA and azide free) version of this product see product #56401. Sitivity TBK1/NAK (E9H5S) Mouse mAb recognizes endogenous levels of total TBK1/NAK protect to sequence Monoclonal antibody is produced by immunizing animals with a synthetic peptide or residues surrounding Glu540 of human TBK1/NAK protein. TBK1 (TANK-binding kinase 1)/NAK (NF-κB activating kinase) is an IκB kinase (IKK)-accan activate IκK through direct phosphorylation (1). TBK1 was identified through as: TRAF binding protein, TANK, and found to function upstream of NIK and IKK in the a (2). TBK1 induces IκB degradation and NF-κB activity through IKKβ. TBK1 may media activation in response to growth factors that stimulate PKCε activity (1). TBK1 plays a activation of IRF3 in the innate immune response (3). Eferences 1. Tojima, Y. et al. (2000) Nature 404, 778-82. 2. Pomerantz, J.L. and Baltimore, D. (1999) EMBO J 18, 6694-704.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting W-F: Fluorescent Western IP: Immunoprecipitation

Cross-Reactivity Key

H: Human M: Mouse R: Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any

purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.