

#5256 Store at -20°C

Phospho-AMPK α (Thr172) (40H9) Rabbit mAb (Biotinylated)



Orders ■ 877-616-CELL (2355)
 orders@cellsignal.com
Support ■ 877-678-TECH (8324)
 info@cellsignal.com
Web ■ www.cellsignal.com

rev. 01/29/16


For Research Use Only. Not For Use In Diagnostic Procedures.

Applications W Endogenous	Species Cross-Reactivity* H, M, R, Hm, Mk, Dm, Sc	Molecular Wt. 62 kDa	Isotype Rabbit IgG
---------------------------------	--	-------------------------	-----------------------


Description: This Cell Signaling Technology antibody is conjugated to biotin under optimal conditions. The biotinylated antibody is expected to exhibit the same species cross-reactivity as the unconjugated Phospho-AMPK α (Thr172) (40H9) Rabbit mAb #2535.

Background: AMP-activated protein kinase (AMPK) is highly conserved from yeast to plants and animals and plays a key role in the regulation of energy homeostasis (1). AMPK is a heterotrimeric complex composed of a catalytic α subunit and regulatory β and γ subunits, each of which is encoded by two or three distinct genes (α 1, 2; β 1, 2; γ 1, 2, 3) (2). The kinase is activated by an elevated AMP/ATP ratio due to cellular and environmental stress, such as heat shock, hypoxia and ischemia (1). The tumor suppressor LKB1, in association with accessory proteins STRAD and MO25, phosphorylates AMPK α at Thr172 in the activation loop and this phosphorylation is required for AMPK activation (3-5). AMPK α is also phosphorylated at Thr258 and Ser485 (for α 1; Ser491 for α 2). The upstream kinase and the biological significance of these phosphorylation events have yet to be elucidated (6). The β 1 subunit is post-translationally modified by myristoylation and multi-site phosphorylation including Ser24/25, Ser96, Ser101, Ser108 and Ser182 (6,7). Phosphorylation at Ser108 of the β 1 subunit seems to be required for the activation of AMPK enzyme, while phosphorylation at Ser24/25 and Ser182 affects AMPK localization (7). Several mutations in AMPK γ subunits have been identified, most of which are located in the putative AMP/ATP binding sites (CBS or Bateman domains). Mutations at these sites lead to reduction of AMPK activity and cause glycogen accumulation in heart or skeletal muscle (1,2). Accumulating evidence indicates that AMPK not only regulates the metabolism of fatty acids and glycogen, but also modulates protein synthesis and cell growth through EF2 and TSC2/mTOR pathways, as well as blood flow via eNOS/nNOS (1).

Specificity/Sensitivity: Phospho-AMPK α (Thr172) (40H9) Rabbit mAb (Biotinylated) detects endogenous AMPK α only when phosphorylated at Thr172. The antibody detects both α 1 and α 2 isoforms of the catalytic subunit, but does not detect the regulatory β or γ subunits.



Western blot analysis of extracts from C2C12 cells, untreated (-) or hydrogen peroxide-treated (+) (AMPK Control Cell Extracts #9158), using Phospho-AMPK α (Thr172) (40H9) Rabbit mAb (Biotinylated).



Western blot analysis of extracts from oligomycin-treated (0.5 μ M) 293 cells either untreated (-) or λ phosphatase (+), using Phospho-AMPK α (Thr172) (40H9) Rabbit mAb (Biotinylated).

Source/Purification: Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Thr172 of human AMPK α protein.

Entrez-Gene ID #5562
UniProt ID #Q13131

Storage: Supplied in 136 mM NaCl, 2.6 mM KCl, 12 mM sodium phosphate (pH 7.4) dibasic, 2 mg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Biotinylated antibodies are designed to be detected using streptavidin conjugates.

***Species cross-reactivity other than human, mouse, monkey and rat is determined by western using the unconjugated antibody.**

Recommended Antibody Dilutions:
 Western blotting 1:1000

For application specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

- Background References:**
- (1) Hardie, D.G. (2004) *J Cell Sci* 117, 5479-87.
 - (2) Carling, D. (2004) *Trends Biochem Sci* 29, 18-24.
 - (3) Hawley, S.A. et al. (1996) *J Biol Chem* 271, 27879-87.
 - (4) Lizcano, J.M. et al. (2004) *EMBO J* 23, 833-43.
 - (5) Shaw, R.J. et al. (2004) *Proc Natl Acad Sci USA* 101, 3329-35.
 - (6) Woods, A. et al. (2003) *J Biol Chem* 278, 28434-42.
 - (7) Warden, S.M. et al. (2001) *Biochem J* 354, 275-83.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween[®]20 at 4°C with gentle shaking, overnight.

© 2014 Cell Signaling Technology, Inc. Cell Signaling Technology[®] is a trademark of Cell Signaling Technology, Inc.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
 Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.

U.S. Patent No. 5,675,063
 Tween[®]20 is a registered trademark of ICI Americas, Inc.