

PDGF Receptor α (D13C6) XP® Rabbit mAb (Biotinylated)



Orders ■ 877-616-CELL (2355)
 orders@cellsignal.com
Support ■ 877-678-TECH (8324)
 info@cellsignal.com
Web ■ www.cellsignal.com

rev. 01/29/16


For Research Use Only. Not For Use In Diagnostic Procedures.

Applications W Endogenous	Species Cross-Reactivity H	Molecular Wt. 190 kDa	Isotype Rabbit IgG
---------------------------------	-------------------------------	--------------------------	-----------------------

Description: This Cell Signaling Technology antibody is conjugated to biotin under optimal conditions. The biotinylated antibody is expected to exhibit the same species cross-reactivity as the unconjugated PDGF Receptor α (D13C6) XP® Rabbit mAb #5241.

Background: The proteins of the platelet derived growth factor (PDGF) family exist as several disulphide-bonded, dimeric isoforms (PDGF AA, PDGF AB, PDGF BB, PDGF CC and PDGF DD) that bind in a specific pattern to two closely related receptor tyrosine kinases, PDGF receptor α (PDGFR α) and PDGF receptor β (PDGFR β). PDGFR α and PDGFR β share 75% to 85% sequence homology between their two intracellular kinase domains while the kinase insert and carboxy-terminal tail regions display a lower level (27% to 28%) of homology (1). PDGF receptor α homodimers bind all PDGF isoforms except those containing PDGF D. PDGF receptor β homodimers bind PDGF BB and DD isoforms, as well as the PDGF AB heterodimer. The heteromeric PDGF α / β receptor binds PDGF B, C, and D homodimers as well as the PDGF AB heterodimer (2). PDGFR α and PDGFR β can each form heterodimers with EGFR, which is also activated by PDGF (3). Various cells differ in the total number of receptors present and in the receptor subunit composition, which may account for responsive differences among cell types to PDGF binding (4). Ligand binding induces receptor dimerization and autophosphorylation, followed by binding and activation of cytoplasmic SH2 domain-containing signal transduction molecules such as Grb2, Src, GAP, PI3 kinase, PLC γ and Nck. A number of different signaling pathways are initiated by activated PDGF receptors and lead to control of cell growth, actin reorganization, migration, and differentiation (5). Tyr751 in the kinase-insert region of PDGFR β is the docking site for PI3 kinase (6). Phosphorylated pentapeptides derived from Tyr751 of PDGFR β (pTyr751-Val-Pro-Met-Leu) inhibit the association of the carboxy-terminal SH2 domain of the p85 subunit of PI3 kinase with PDGFR β (7). Tyr740 is also required for PDGFR β -mediated PI3 kinase activation (8).

Specificity/Sensitivity: PDGF Receptor α (D13C6) XP® Rabbit mAb (Biotinylated) recognizes endogenous levels of total PDGF Receptor α protein.



Western blot analysis of extracts from NCI-H1703, HCC827, and U-118 MG cells using PDGF Receptor α (D13C6) XP® Rabbit mAb (Biotinylated).

Source/Purification: Monoclonal antibody is produced by immunizing animals with a recombinant protein corresponding to the PDGF receptor α extracellular domain.

Entrez-Gene ID #5156
 Swiss-Prot Acc. #P16234

Storage: Supplied in 136 mM NaCl, 2.6 mM KCl, 12 mM sodium phosphate (pH 7.4) dibasic, 2 mg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot antibody.

Biotinylated antibodies are designed to be detected using streptavidin or anti-Biotin antibody conjugates.

Recommended Antibody Dilutions:
 Western Blotting 1:1000

For application specific protocols please see the web page for this product at www.cellsignal.com.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

Background References:

- (1) Deuel, T.F. et al. (1988) *Biofactors* 1, 213-217.
- (2) Bergsten, E. et al. (2001) *Nat. Cell Biol.* 3, 512-516.
- (3) Betscholtz, C. et al. (2001) *Bioessays* 23, 494-507.
- (4) Coughlin, S.R. et al. (1988) *Prog. Clin. Biol. Res.* 266, 39-45.
- (5) Ostman, A. and Heldin, C.H. (2001) *Adv. Cancer Res.* 80, 1-38.
- (6) Panayotou, G. et al. (1992) *EMBO J.* 11, 4261-4272.
- (7) Ramalingam, K. et al. (1995) *Bioorg. Med. Chem.* 3, 1263-1272.
- (8) Kashishian, A. et al. (1992) *EMBO J.* 11, 1373-1382.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
 Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.