

2B4/SLAMF4/CD244 (D5J9D) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H M	Endogenous	70-120	Rabbit IgG	#Q9BZW8	51744

Product Usage Information**Application**

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:100

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. *Do not aliquot the antibody.*

Specificity/Sensitivity

2B4/SLAMF4/CD244 (D5J9D) Rabbit mAb recognizes endogenous levels of total 2B4/SLAMF4/CD244 protein. This antibody cross-reacts with an unidentified protein of 18 kDa in some cell extracts.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Asp367 of human 2B4/SLAMF4/CD244 protein.

Background

2B4, also known as signaling lymphocytic activation molecule family member 4 (SLAMF4) and cluster of differentiation 244 (CD244), is a heterophilic cell surface receptor expressed on a variety of immune cells, including natural killer (NK) cells, T cells, eosinophils, mast cells, and dendritic cells. 2B4 has been shown to have both immune stimulatory and inhibitory effects on cells (1-11). Upon engagement of its ligand CD48, 2B4 can enhance immune cell signaling, cytokine production, non-major histocompatibility complex-restricted cytotoxicity, and cell migration (1-8). Conversely, at early stages of NK cell differentiation, 2B4 may function as an inhibitory receptor possibly ensuring the self-tolerance of developing NK cells (10). 2B4 also inhibits inflammatory responses in dendritic cells (11). Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Background References

1. Speiser, D.E. et al. (2001) *J Immunol* 167, 6165-70.
2. Nakajima, H. et al. (1999) *Eur J Immunol* 29, 1676-83.
3. Valiante, N.M. and Trinchieri, G. (1993) *J Exp Med* 178, 1397-406.
4. Munitz, A. et al. (2005) *J Immunol* 174, 110-8.
5. Sivori, S. et al. (2000) *Eur J Immunol* 30, 787-93.
6. El-Shazly, A.E. et al. *Int J Immunopathol Pharmacol* 24, 949-60.
7. Chen, L. and Flies, D.B. (2013) *Nat Rev Immunol* 13, 227-42.
8. Assarsson, E. et al. (2005) *Mol Immunol* 42, 419-23.
9. Elishmereni, M. et al. (2014) *J Invest Dermatol* 134, 2928-37.
10. Sivori, S. et al. (2002) *Proc Natl Acad Sci U S A* 99, 4526-31.
11. Georgoudaki, A.M. et al. (2015) *Immunol Cell Biol* 93, 581-90.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.