#5513 Store at -20C

Estrogen Receptor β Antibody

Cell Signaling TECHNOLOGY®

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source:	UniProt ID:	Entrez-Gene Id:
WB	H M R Mk	Endogenous	52, 55, 63	Rabbit	#Q92731	2100

Product Usage Information

Application	Dilution
Western Blotting	1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at –20°C. Do not aliquot the antibody.

Specificity / Sensitivity

Estrogen Receptor β Antibody detects endogenous levels of total Estrogen Receptor β protein. This antibody is predicted to cross-react with all Estrogen Receptor β isoforms. This antibody does not cross-react with Estrogen Receptor α.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the amino terminus of human Estrogen Receptor β1 protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

Estrogen Receptor β (ER β) is a member of the nuclear receptor superfamily of transcription factors and was discovered to be encoded by a gene (*ESR2)* distinct from that encoding Estrogen Receptor α (ER α) (1,2). While studies have revealed that alternative splicing generates mutiple isoforms of ER β that differ at their respective C-termini and in tissue distribution, ER β1 is believed to be the longest and only fully functional isoform (3,4). Indeed, it has been reported that shorter isoforms of ER β (ER β2, β4, and β5) can heterodimerize with ER β1 and enhance its transcriptional activity in an estradiol-dependent manner (4). ER β is expressed in a wide range of normal and malignant tissues, many of which coexpress ER α. It is proposed that ER β has an antiproliferative role, perhaps through heterodimerization with ER α and repression of its transcriptional activity at estrogen response elements (5,6). Recent studies have revealed that expression of *ESR2* is subject to epigenetic regulation and that loss of ER β expression positively contributes to epithelial-mesenchymal transition and enhanced invasiveness in prostate cancer (7,8). ER β has also been found to be negatively regulated at the posttranslational level through phosphorylation of its AF-1 domain, which promotes its ubiquitin-dependent proteasomal degradation (9,10).

Background References

1. Evans, R.M. (1988) *Science* 240, 889-95.
2. Kuiper, G.G. et al. (1996) *Proc Natl Acad Sci U S A* 93, 5925-30.
3. Moore, J.T. et al. (1998) *Biochem Biophys Res Commun* 247, 75-8.
4. Leung, Y.K. et al. (2006) *Proc Natl Acad Sci U S A* 103, 13162-7.
5. Hall, J.M. and McDonnell, D.P. (1999) *Endocrinology* 140, 5566-78.
6. Pettersson, K. et al. (2000) *Oncogene* 19, 4970-8.
7. Zhu, X. et al. (2004) *Am J Pathol* 164, 2003-12.
8. Mak, P. et al. (2010) *Cancer Cell* 17, 319-32.
9. Tateishi, Y. et al. (2006) *Mol Cell Biol* 26, 7966-76.
10. Picard, N. et al. (2008) *Mol Endocrinol* 22, 317-30.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

WB: Western Blotting

Cross-Reactivity Key

H: human **M:** mouse **R:** rat **Hm:** hamster **Mk:** monkey **Vir:** virus **Mi:** mink **C:** chicken **Dm:** D. melanogaster **X:** Xenopus **Z:** zebrafish **B:** bovine **Dg:** dog **Pg:** pig **Sc:** S. cerevisiae **Ce:** C. elegans **Hr:** horse **GP:** Guinea Pig **Rab:** rabbit **All:** all species expected

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.
All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.