

#5537 Store at -20C

HIF-1 β /ARNT (D28F3) XP[®] Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP, IHC-P, ChIP, ChIP-seq, C&R	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 87	Source/Isotype: Rabbit IgG	UniProt ID: #P27540	Entrez-Gene Id: 405
--	--------------------------------	-----------------------------------	------------------------	--------------------------------------	-------------------------------	-------------------------------

Product Usage Information

For optimal ChIP and ChIP-seq results, use 10 μ l of antibody and 10 μ g of chromatin (approximately 4 x 10⁶ cells) per IP. This antibody has been validated using SimpleChIP[®] Enzymatic Chromatin IP Kits.

The CUT&RUN dilution was determined using CUT&RUN Assay Kit #86652.

Application	Dilution
Western Blotting	1:1000
Immunoprecipitation	1:50
Immunohistochemistry (Paraffin)	1:50 - 1:200
Chromatin IP	1:50
Chromatin IP-seq	1:50
CUT&RUN	1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

For a carrier free (BSA and azide free) version of this product see product #43589.

Specificity/Sensitivity

HIF-1 β /ARNT (D28F3) XP[®] Rabbit mAb detects endogenous levels of total HIF-1 β /ARNT protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to the sequence around Ile479 of human HIF-1 β /ARNT protein.

Background

Hypoxia-inducible factor 1 (HIF1) is a heterodimeric transcription factor that plays a critical role in the cellular response to hypoxia (1). The HIF1 complex consists of two subunits, HIF-1 α and HIF-1 β , which are basic helix-loop-helix proteins of the PAS (Per, ARNT, Sim) family (2). HIF1 regulates the transcription of a broad range of genes that facilitate responses to the hypoxic environment, including genes regulating angiogenesis, erythropoiesis, cell cycle, metabolism, and apoptosis. The widely expressed HIF-1 α is typically degraded rapidly in normoxic cells by the ubiquitin/proteasomal pathway. Under normoxic conditions, HIF-1 α is proline hydroxylated leading to a conformational change that promotes binding to the von Hippel-Lindau protein (VHL) E3 ligase complex; ubiquitination and proteasomal degradation follows (3,4). Both hypoxic conditions and chemical hydroxylase inhibitors (such as desferrioxamine and cobalt) inhibit HIF-1 α degradation and lead to its stabilization. In addition, HIF-1 α can be induced in an oxygen-independent manner by various cytokines through the PI3K-AKT-mTOR pathway (5-7).

HIF-1 β is also known as AhR nuclear translocator (ARNT) due to its ability to partner with the aryl hydrocarbon receptor (AhR) to form a heterodimeric transcription factor complex (8). Together with AhR, HIF-1 β plays an important role in xenobiotics metabolism (8). In addition, a chromosomal translocation leading to a TEL-ARNT fusion protein is associated with acute myeloblastic leukemia (9). Studies also found that ARNT/HIF-1 β expression levels decrease significantly in pancreatic islets from patients with type 2 diabetes, suggesting that HIF-1 β plays an important role in pancreatic β -cell function (10).

Background References

1. Sharp, F.R. and Bernaudin, M. (2004) *Nat Rev Neurosci* 5, 437-48.
 2. Wang, G.L. et al. (1995) *Proc Natl Acad Sci U S A* 92, 5510-4.
 3. Jaakkola, P. et al. (2001) *Science* 292, 468-72.
 4. Maxwell, P.H. et al. (1999) *Nature* 399, 271-5.
 5. Fukuda, R. et al. (2002) *J Biol Chem* 277, 38205-11.
 6. Jiang, B.H. et al. (2001) *Cell Growth Differ* 12, 363-9.
 7. Laughner, E. et al. (2001) *Mol Cell Biol* 21, 3995-4004.
 8. Walisser, J.A. et al. (2004) *Proc Natl Acad Sci U S A* 101, 16677-82.
 9. Salomon-Nguyen, F. et al. (2000) *Proc Natl Acad Sci U S A* 97, 6757-62.
 10. Gunton, J.E. et al. (2005) *Cell* 122, 337-49.
-

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation **IHC-P:** Immunohistochemistry (Paraffin) **ChIP:** Chromatin IP **ChIP-seq:** Chromatin IP-seq **C&R:** CUT&RUN

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.