909

BACE1 (D10E5) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only, Not for Use in Diagnostic Procedures

Applications: W, IP, IF-F, IF-IC	Reactivity: H M R	Sensitivity: Endogenous	MW (kDa): 70	Source/Isotype: Rabbit IgG	UniProt ID: #P56817	Entrez-Gene Id 23621
Product Usage Information		Application			Dilution	
		Western Blotting			1:1	000
		Immunoprecipitation	1		1:5	0
		Immunofluorescence	e (Frozen)		1:5	0 - 1:200
		Immunofluorescence	(Immunocytochem	istry)	1:2	00 - 1:800
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
		For a carrier-free (BSA and azide free) version of this product see product #20562.				
Specificity/Sensitivity		BACE1 (D10E5) Rabbit mAb detects endogenous levels of total BACE1 protein. This product does not detect BACE2.				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding His490 of human BACE1 protein.				
Background		β-secretase (BACE) is an aspartic acid proteinase that catalyzes the initial step in APP processing by cleaving and releasing a soluble, extracellular APP- $β$ (sAPP $β$) ectodomain and generating a membrane-bound, carboxy-terminal fragment consisting of 99 amino acids (CTF99). Additional processing of CTF99 by $γ$ -secretase generates the amyloid $β$ -peptide ($Aβ$) that forms aggregates in the brains of Alzheimer disease (AD) patients. BACE is an attractive target for inhibitors in AD therapy since it catalyzes the first and rate limiting step in amyloidogenic APP processing (1). Pro-BACE-1 is synthesized in the ER before it is transported to the trans-Golgi network to undergo maturation (2). BACE-1 functions as a dimer and is phosphorylated at Ser498 by casein kinase 1, which is necessary for recycling back to the membrane from the early endosome (3,4).				
Background References		1. Hunt, C.E. and Turner, A.J. (2009) <i>FEBS J</i> 276, 1845-59. 2. Benjannet, S. et al. (2001) <i>J Biol Chem</i> 276, 10879-87. 3. Walter, J. et al. (2001) <i>J Biol Chem</i> 276, 14634-41. 4. He, X. et al. (2005) <i>J Biol Chem</i> 280, 11696-703.				
Species Reacti	vity	Species reactivity is d	etermined by testin	g in at least one approve	ed application (e.g.,	western blot).
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.				

Applications Key W: Western Blotting IP: Immunoprecipitation IF-F: Immunofluorescence (Frozen) IF-IC:

Immunofluorescence (Immunocytochemistry)

Cross-Reactivity Key H: Human M: Mouse R: Rat

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for

more information.

Limited UsesExcept as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.