## Phospho-SMAD2 (Ser465/Ser467) (E8F3R) Rabbit mAb (Alexa Fluor® 488 Conjugate)



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

| <b>Applications:</b><br>FC-FP | Reactivity:<br>H M R | <b>Sensitivity:</b><br>Endogenous  | <b>Source/Isotype:</b><br>Rabbit IgG | UniProt ID:<br>#Q15796 | Entrez-Gene Id:<br>4087             |
|-------------------------------|----------------------|--|--------------------------------------|------------------------|-------------------------------------|
| Product Usage<br>Information  |                      | <b>Application</b> Flow Cytometry (Fixed/P   | ermeabilized)                        |                        | <b>Dilution</b><br>1:50             |
| Storage                       |                      | Supplied in PBS (pH 7.2), antibody. Protect from lig   |                                      | zide and 2 mg/ml BS    | A. Store at 4°C. Do not aliquot the |
| Specificity/Sensitivity       |                      | Phospho-Smad2 (Ser465/Ser467) (E8F3R) Rabbit mAb (Alexa Fluor <sup>®</sup> 488 Conjugate) recognizes endogenous levels of Smad2 protein when phosphorylated at Ser465 and Ser467.  |                                      |                        |                                     |
| Source / Purification         |                      | Monoclonal antibody is produced by immunizing animals with a synthetic phosphopeptide corresponding to residues surrounding Ser465/Ser467 of human Smad2 protein.  |                                      |                        |                                     |
| Description                   |                      | This Cell Signaling Technology antibody is conjugated to Alexa Fluor <sup>®</sup> 488 fluorescent dye and tested in-house for direct flow cytometric analysis in human cells. This antibody is expected to exhibit the same species cross-reactivity as the unconjugated Phospho-Smad2 (Ser465/Ser467) (E8F3R) Rabbit mAb #18338.  |                                      |                        |                                     |
| Background                    |                      | Members of the SMAD family of signal transduction molecules are components of a critical intracellular pathway that transmit TGF- $\beta$ signals from the cell surface into the nucleus. Three distinct classes of SMADs have been defined: the receptor-regulated SMADs (R-SMADs), which include SMAD1, 2, 3, 5, and 9; the common-mediator SMAD (co-SMAD), SMAD4; and the antagonistic or inhibitory SMADs (I-SMADs), SMAD6 and 7 (1-5). Activated type I receptors associate with specific R-SMADs and phosphorylate them on a conserved carboxy-terminal SSXS motif. The phosphorylated R-SMADs dissociate from the receptor and form a heteromeric complex with SMAD4, initiating translocation of the heteromeric SMAD complex to the nucleus. Once in the nucleus, SMADs recruit a variety of DNA binding proteins that function to regulate transcriptional activity (6-8). |                                      |                        |                                     |
| Background References         |                      | <ol> <li>Heldin, C.H. et al. (1997) Nature 390, 465-71.</li> <li>Attisano, L. and Wrana, J.L. (1998) Curr Opin Cell Biol 10, 188-94.</li> <li>Derynck, R. et al. (1998) Cell 95, 737-40.</li> <li>Massagué, J. (1998) Annu Rev Biochem 67, 753-91.</li> <li>Whitman, M. (1998) Genes Dev 12, 2445-62.</li> <li>Wrana, J.L. (2000) Sci STKE 2000, re1.</li> <li>Attisano, L. and Wrana, J.L. (2002) Science 296, 1646-7.</li> <li>Moustakas, A. et al. (2001) J Cell Sci 114, 4359-69.</li> </ol>   |                                      |                        |                                     |
|                               |                      |  |                                      |                        |                                     |

**Species Reactivity** 

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Applications Key** FC-FP: Flow Cytometry (Fixed/Permeabilized)

Cross-Reactivity Key H: Human M: Mouse R: Rat

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

This product is provided under an intellectual property license from Life Technologies Corporation. The transfer of this product is conditioned on the buyer using the purchased product solely in research conducted by the buyer, excluding contract research or any fee for service research, and the buyer must not (1) use this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; or (c) manufacturing or quality assurance or quality control, and/or (2) sell or transfer this product or its components for resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life

Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

## **Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.