



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

RMP Antibody

Store at -20C
#5844

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: H M R Mk	Sensitivity: Endogenous	MW (kDa): 79	Source/Isotype: Rabbit	UniProt ID: #O94763	Entrez-Gene Id: 8725
-------------------------------	--------------------------------	-----------------------------------	------------------------	----------------------------------	-------------------------------	--------------------------------

Product Usage Information

Application

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA and 50% glycerol. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

RMP Antibody recognizes endogenous levels of total RMP protein.

Source / Purification

Polyclonal antibodies are produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human RMP protein. Antibodies are purified by protein A and peptide affinity chromatography.

Background

RMP (RBP5-Mediating Protein), also known as URI (Unconventional prefoldin RBP5 Interactor), was described as an unconventional member of the prefoldin (PFD) family of chaperones that are involved in actin and tubulin folding (1-4). Like conventional members of the α -class of PFDs, RMP contains N- and C-terminal α -helical coiled-coil structures connected by two β hairpins. In addition, RMP possesses an RBP5-binding segment and a long C-terminal acidic segment. It is posited that RMP exists as a component of a macromolecular complex within human cells and functions as a molecular scaffold to assemble a PFD complex containing other PFDs and proteins with functions in transcription and ubiquitination. Indeed, evidence is provided that RMP negatively modulates RNA polymerase II-dependent transcription by binding to TFIIF (5) and RBP5 (6) and is involved in mTOR signaling by coordinating the regulation of nutrient availability with gene expression (1). In accord with its ability to coordinate gene expression with nutrient availability, RMP was shown to be a mitochondrial substrate of S6K1. S6K1-mediated phosphorylation of RMP at Ser371 triggers a series of biochemical events that constitute a negative feedback loop, in part, aimed at restraining S6K1 survival signaling and ensuring that the mitochondrial threshold for apoptosis corresponds to availability of nutrients and growth factors (7).

Background References

- Gstaiger, M. et al. (2003) *Science* 302, 1208-12.
- Vainberg, I.E. et al. (1998) *Cell* 93, 863-73.
- Martín-Benito, J. et al. (2002) *EMBO J* 21, 6377-86.
- Geissler, S. et al. (1998) *EMBO J* 17, 952-66.
- Wei, W. et al. (2003) *Cell Res* 13, 111-20.
- Dorjsuren, D. et al. (1998) *Mol Cell Biol* 18, 7546-55.
- Djouder, N. et al. (2007) *Mol Cell* 28, 28-40.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat **Mk:** Monkey

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.