



Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

Store at -20C
#5864

CA12 (D75C6) XP[®] Rabbit mAb

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP, FC-L	Reactivity: H	Sensitivity: Endogenous	MW (kDa): 45-50	Source/Isotype: Rabbit IgG	UniProt ID: #O43570	Entrez-Gene Id: 771
-------------------------------------	-------------------------	-----------------------------------	---------------------------	--------------------------------------	-------------------------------	-------------------------------

Product Usage Information

Application

Western Blotting
Immunoprecipitation
Flow Cytometry (Live)

Dilution

1:1000
1:200
1:200

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

CA12 (D75C6) XP[®] Rabbit mAb detects endogenous levels of total CA12 protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly39 of human CA12 protein.

Background

Carbonic anhydrases (CA) are a family of ancient zinc metalloenzymes found in almost all living organisms. All CA can be divided into 3 distinct classes (α, β, and γ) that evolved independently and have no significant homology in sequence and overall folding. All functional CA catalyze the reversible hydration of CO₂ into HCO₃⁻ and H⁺ and contain a zinc atom in the active sites essential for catalysis. There are many isoforms of CA in mammals and they all belong to the α class (1,2). CA12, a member of the α class, is a plasma membrane protein with the catalytic domain in the extracellular space. CA12 is widely expressed in many tissues and is found at high levels in renal cell cancers. Its expression is induced by hypoxia in solid tumors and is considered to be a tumor marker (3-5). Genetic mutations in CA12 are associated with hyperchlorhidrosis (6,7).

Background References

1. Smith, K.S. et al. (1999) *Proc Natl Acad Sci USA* 96, 15184-9.
2. Tripp, B.C. et al. (2001) *J Biol Chem* 276, 48615-8.
3. Chiche, J. et al. (2009) *Cancer Res* 69, 358-68.
4. Barnett, D.H. et al. (2008) *Cancer Res* 68, 3505-15.
5. Watson, P.H. et al. (2003) *Br J Cancer* 88, 1065-70.
6. Feldshtein, M. et al. (2010) *Am J Hum Genet* 87, 713-20.
7. Muhammad, E. et al. (2011) *Hum Genet* 129, 397-405.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation **FC-L:** Flow Cytometry (Live)

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.