# BRSK1 (D10F2) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

## For Research Use Only. Not for Use in Diagnostic Procedures.

<b>Applications:</b> W, IP	<b>Reactivity:</b> H M R	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 90	<b>Source/Isotype:</b> Rabbit IgG	UniProt ID: #Q8TDC3	Entrez-Gene Id: 84446
Product Usage Information		Application Western Blotting Immunoprecipitation			<b>Dilution</b> 1:1000 1:50	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 $\mu$ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		BRSK1 (D10F2) Rabbit mAb detects endogenous levels of total BRSK1 protein.				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Gly485 of human BRSK1 protein.				
Background		<b>Br</b> ain- <b>s</b> pecific <b>k</b> inase <b>1</b> (BRSK1; SAD-B) and <b>Br</b> ain- <b>s</b> pecific <b>k</b> inase 2 (BRSK2; SAD-A) are serine/threonine kinases closely related to AMPK. LKB1 phosphorylates Thr189 in the T-loop of BRSK1 and Thr174 in the T-loop of BRSK2, resulting in activation of the kinases (1). BRSK1 localizes to synaptic vesicles in the hippocampus and cerebellum, together with the active zone proteins Bassoon and CAST, and BRSK1 phoshorylates the active zone protein RIM1 (2). An alternatively spliced from of BRSK1 displays unique activity during the cell cycle, phosphorylating Ser131 of γ-tubulin and controling centrosome duplication (3). Neuronal polarization, including axon formation, is fundamental for normal brain development. BRSK1 -/- and BRSK2 -/- mice have defects in neuronal polarity and impaired corticogenesis (4). Knockdown of BRSK1 and BRSK2 <i>in vitro</i> diminishes axonal growth (5).				
Background References		<ol> <li>Bright, N.J. et al. (2008) J Biol Chem 283, 14946-54.</li> <li>Inoue, E. et al. (2006) Neuron 50, 261-75.</li> <li>Alvarado-Kristensson, M. et al. (2009) Nat Cell Biol 11, 1081-92.</li> <li>Kishi, M. et al. (2005) Science 307, 929-32.</li> <li>Choi, Y.J. et al. (2008) Genes Dev 22, 2485-95.</li> </ol>				
Species Reactivity		Species reactivity is determined by testing in at least one approved application (e.g., western blot).				
Western Blot Buffer		IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X				

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

# Applications Key

W: Western Blotting IP: Immunoprecipitation

# Cross-Reactivity Key

H: Human M: Mouse R: Rat

#### **Trademarks and Patents**

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

### **Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for

diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.