

Store at
-20°C

Ghost Dye™ Violet 510 Viability Dye

Cell Signaling
TECHNOLOGY®

#59863



100 µl

Support: +1-978-867-2388 (U.S.)
www.cellsignal.com/supportOrders: 877-616-2355 (U.S.)
orders@cellsignal.com

rev. 03/27/20

For Research Use Only. Not For Use In Diagnostic Procedures.Applications
FSpecies Cross-Reactivity
All

Description: Ghost Dye™ Violet 510 Viability Dye is used to discriminate viable from non-viable mammalian cells in flow cytometry applications. Ghost Dye™ Violet 510 Viability Dye irreversibly binds free amines available on the cell surface as well as intracellular free amines exposed in cells with compromised cell membranes. Non-viable cells with loss of membrane integrity will react with significantly more Ghost Dye™ Violet 510 Viability Dye than healthy cells in the same sample. Cells that exhibit increased fluorescence intensity can be excluded from analysis.



Flow cytometric analysis of live and fixed/permeabilized human peripheral blood mononuclear cells, combined and stained with Ghost Dye™ Violet 510 Viability Dye. Viable gate is indicated.

Storage: Store at -20°C desiccated and protected from light. This product is stable for 12 months. Aliquot to avoid excessive freeze-thaw cycles.

Directions For Use:

1. Prepare the following reagents with reverse osmosis deionized (RODI) or equivalent grade water:
 - a. 1X PBS (azide- and protein/serum-free)
 - b. Incubation Buffer: Dissolve 0.5 g Bovine Serum Albumin (BSA) (#9998) in 100 ml 1X PBS. Store at 4°C.
2. Remove Ghost Dye™ from -20°C and bring to room temperature.
3. Collect cells by centrifugation and aspirate supernatant.
4. Wash cells by centrifugation in excess 1X PBS. Repeat if necessary.
5. Resuspend cells to a concentration of 1-10 x 10⁶/mL in 1X PBS.
6. Centrifuge the Ghost Dye™ before opening then add 1 µL for each 1 mL of cell suspension and vortex immediately.
7. Incubate for 30 minutes at 4°C protected from light.
8. Wash by centrifugation in excess incubation buffer. Discard supernatant. Repeat.
9. Cells can then be stained, fixed and/or permeabilized based upon experimental design.

Ghost Dye™ Violet 510 Viability Dye is excited by the violet (405 nm) laser line and has a peak emission of 510 nm that can be detected using a 525/50 band pass filter commonly used for detection of AmCyan.

Ghost Dye is a registered trademark of Tonbo Biosciences.

Thank you for your recent purchase. If you would like to provide a review visit cellsignal.com/comments.

www.cellsignal.com

© 2017 Cell Signaling Technology, Inc.

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Applications: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide Species Cross-Reactivity: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.