

#6235 Store at -20°C

SignalSilence® Jak2 siRNA I



✓ 10 µM in 300 µl (100 transfections)

Orders ■ 877-616-CELL (2355) orders@cellsignal.com
Support ■ 877-678-TECH (8324) info@cellsignal.com
Web ■ www.cellsignal.com

rev. 03/08/16

For Research Use Only. Not For Use In Diagnostic Procedures.


Species Cross-Reactivity: H

Description: SignalSilence® Jak2 siRNA I from Cell Signaling Technology (CST) allows the researcher to specifically inhibit Jak2 expression using RNA interference, a method whereby gene expression can be selectively silenced through the delivery of double stranded RNA molecules into the cell. All SignalSilence® siRNA products from CST are rigorously tested in-house and have been shown to reduce target protein expression by western analysis.

Background: Members of the Janus family of tyrosine kinases (Jak1, Jak2, Jak3 and Tyk2) are activated by ligands binding to a number of associated cytokine receptors (1). Upon cytokine receptor activation, Jak proteins become autophosphorylated and phosphorylate their associated receptors to provide multiple binding sites for signaling proteins. These associated signaling proteins, such as Stats (2), Shc (3), insulin receptor substrates (4) and focal adhesion kinase (FAK) (5), typically contain SH2 or other phospho-tyrosine-binding domains.

Directions for Use: CST recommends transfection with 100 nM Jak2 siRNA I 48 to 72 hours prior to cell lysis. For transfection procedure, follow protocol provided by the transfection reagent manufacturer. Please feel free to contact CST with any questions on use.

Quality Control: Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid phase extraction. The annealed RNA duplex is further analyzed by mass spectrometry to verify the exact composition of the duplex. Each lot is compared to the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.



Western blot analysis of extracts from K-562 cells, transfected with 100 nM SignalSilence® Control siRNA (Unconjugated) #6568 (-) or SignalSilence® Jak2 siRNA I (+), using Jak2 (D2E12) XP™ Rabbit mAb #3230 and α-Tubulin (11H10) Rabbit mAb #2125. The Jak2 (D2E12) XP™ Rabbit mAb confirms silencing of Jak2 expression, while the α-Tubulin (11H10) Rabbit mAb is used to control for loading and specificity of Jak2 siRNA.

Entrez-Gene ID #3717
Swiss-Prot Acc. #060674

Storage: Jak2 siRNA I is supplied in RNase-free water. Aliquot and store at -20°C.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

Background References:

- (1) Leonard, W.J. and O'Shea, J.J. (1998) *Annu. Rev. Immunol.* 16, 293-322.
- (2) Darnell, J.E. (1997) *Science* 277, 1630-1635.
- (3) VanderKuur, J. et al. (1995) *J. Biol. Chem.* 270, 7587-7593.
- (4) Argetsinger, L.S. et al. (1995) *J. Biol. Chem.* 270, 14685-14692.
- (5) Zhu, T. et al. (1998) *J. Biol. Chem.* 273, 10682-10689.

© 2010 Cell Signaling Technology, Inc.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.