

Store at
-20°C

#62849

Na,K-ATPase β 1 (D8W8J) Rabbit mAb

Support: +1-978-867-2388 (U.S.)
www.cellsignal.com/supportOrders: 877-616-2355 (U.S.)
orders@cellsignal.comEntrez-Gene ID #481
UniProt ID #P05026

New 04/17

For Research Use Only. Not For Use In Diagnostic Procedures.

Applications W Endogenous	Species Cross-Reactivity* H, M, R	Molecular Wt. 45-55 kDa	Isotype Rabbit IgG**
---------------------------------	--------------------------------------	----------------------------	-------------------------

Background: The Na,K-ATPase is an integral membrane heterodimer belonging to the P-type ATPase family. This ion channel uses the energy derived from ATP hydrolysis to maintain membrane potential by driving sodium export and potassium import across the plasma membrane against their electrochemical gradients. It is composed of a catalytic α subunit and a β subunit (reviewed in 1). Several phosphorylation sites have been identified for the α 1 subunit. Tyr10 is phosphorylated by an as yet undetermined kinase (2), Ser16 and Ser23 are phosphorylated by PKC, and Ser943 is phosphorylated by PKA (3-5). All of these sites have been implicated in the regulation of enzyme activity in response to hormones and neurotransmitters, altering trafficking and kinetic properties of Na,K-ATPase. Altered phosphorylation in response to angiotensin II stimulates activity in the rat proximal tubule (6). Na,K-ATPase is also involved in other signal transduction pathways. Insulin regulates its localization in differentiated primary human skeletal muscle cells, and this regulation is dependent on ERK1/2 phosphorylation of the α subunit (7). Na,K-ATPase and Src form a signaling receptor complex that affects regulation of Src kinase activity and, subsequently, its downstream effectors (8,9).

Na,K-ATPase β 1 is the non-catalytic subunit of Na,K-ATPase. It is required for stabilization, maturation, and translocation of the catalytic α subunit to the plasma membrane (10-12). Na,K-ATPase β 1 also mediates the trans-dimerization of Na,K-ATPase between neighboring cells where it regulates the integrity of tight junctions (13-17). Glutathionylation of Na,K-ATPase β 1 regulates the ion pump activity of Na,K-ATPase (18). Research studies have shown that Na,K-ATPase β 1 is a target of the Sonic Hedgehog signaling pathway and may be involved in suppressing tumor development and progression (19). *ATP1B1*, the gene encoding Na,K-ATPase β 1, is epigenetically silenced by promoter methylation in both renal cell carcinoma cell lines and patient tissues (20).

Specificity/Sensitivity: Na,K-ATPase β 1 (D8W8J) Rabbit mAb recognizes endogenous levels of total Na,K-ATPase β 1 protein.

Source/Purification: Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Lys85 of human Na,K-ATPase β 1 protein.

Background References:

- (1) Therien, A.G. and Blostein, R. (2000) *Am J Physiol Cell Physiol* 279, C541-66.
- (2) Féraille, E. et al. (1999) *Mol Biol Cell* 10, 2847-59.
- (3) Fisone, G. et al. (1994) *J Biol Chem* 269, 9368-73.
- (4) Feschenko, M.S. and Sweadner, K.J. (1995) *J Biol Chem* 270, 14072-7.
- (5) Beguin, P. et al. (1994) *J Biol Chem* 269, 24437-45.
- (6) Yingst, D.R. et al. (2004) *Am J Physiol Renal Physiol* 287, F713-21.
- (7) Al-Khalili, L. et al. (2004) *J Biol Chem* 279, 25211-8.
- (8) Tian, J. et al. (2006) *Mol Biol Cell* 17, 317-26.
- (9) Liang, M. et al. (2006) *J Biol Chem* 281, 19709-19.
- (10) Beggah, A.T. et al. (1997) *J Biol Chem* 272, 10318-26.
- (11) Hasler, U. et al. (1998) *J Biol Chem* 273, 30826-35.
- (12) Rajasekaran, S.A. et al. (2004) *Mol Biol Cell* 15, 3224-32.
- (13) Rajasekaran, S.A. et al. (2001) *Mol Biol Cell* 12, 3717-32.
- (14) Rajasekaran, A.K. and Rajasekaran, S.A. (2003) *Am J Physiol Renal Physiol* 285, F388-96.
- (15) Bab-Dinitz, E. et al. (2009) *Biochemistry* 48, 8684-91.
- (16) Tokhtaeva, E. et al. (2011) *J Biol Chem* 286, 25801-12.
- (17) Vagin, O. et al. (2012) *Am J Physiol Cell Physiol* 302, C1271-81.
- (18) Figtree, G.A. et al. (2012) *Free Radic Biol Med* 53, 2263-8.
- (19) Lee, S.J. et al. (2015) *Mol Cancer* 14, 159.
- (20) Selvakumar, P. et al. (2014) *Epigenetics* 9, 579-86.

Storage: Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μ g/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.


*Species cross-reactivity is determined by western blot.

**Anti-rabbit secondary antibodies must be used to detect this antibody.

Recommended Antibody Dilutions:

Western blotting 1:1000

For product specific protocols and a complete listing of recommended companion products please see the product web page at www.cellsignal.com



Western blot analysis of extracts from human, mouse, and rat brains using Na,K-ATPase β 1 (D8W8J) Rabbit mAb.

Tween is a registered trademark of ICI Americas, Inc.

Thank you for your recent purchase. If you would like to provide a review visit cellsignal.com/comments.

www.cellsignal.com

© 2017 Cell Signaling Technology, Inc.

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

Applications: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide Species Cross-Reactivity: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.