

## 63182

## UBE2G2 (D8Z4G) Rabbit mAb



Orders: 877-616-CELL (2355)

orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W	Reactivity: H M R Mk	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 19	<b>Source/Isotype:</b> Rabbit IgG	UniProt ID: #P60604	Entrez-Gene Io 7327
Product Usage Information		<b>Application</b> Western Blotting			<b>Dilution</b> 1:1000	
Storage		Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 μg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at –20°C. Do not aliquot the antibody.				
Specificity/Sensitivity		UBE2G2 (D8Z4G) Rabbit mAb recognizes endogenous levels of total UBE2G2 protein. This antibody does not cross-react with UBE2G1 protein.				
Species predicte based on 100% s homology		Bovine, Pig				
Source / Purification		Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human UBE2G2 protein.				
Background		Ubiquitin (Ub) is a conserved polypeptide that is covalently linked to many cellular proteins through the process of ubiquitination, which targets proteins for degradation by the 26S proteasome. Three enzymatic components are involved in the protein ubiquitination cascade. Ubiquitin is first activated by forming a thioester complex with an E1 ubiquitin-activating enzyme. Activated ubiquitin is subsequently transferred to an E2 ubiquitin-carrier protein, and then from the E2 to an E3 ubiquitin ligase for final delivery to the ε-amino group of the target protein lysine residue (1-3). The ubiquitin-conjugating enzyme E2 G2 (UBE2G2, UBC7) is a ubiquitously expressed E2 enzyme and critical component of the endoplasmic reticulum-associated degradation pathway (ERAD) (4). Research studies demonstrate that UBE2G2 forms homodimers and preassembles K48-linked poly-Ub chains at its active site (5-8). The association of Ub-charged UBE2G2 molecules with the ER-resident E3 ligase AMFR (gp78) is required for Ub chain transfer and efficient removal of misfolded or aggregated proteins through the ERAD pathway (9,10).				
Background References		1. Ciechanover, A. (1998) <i>EMBO J</i> 17, 7151-60. 2. Hochstrasser, M. (2000) <i>Nat Cell Biol</i> 2, E153-7. 3. Hochstrasser, M. (2000) <i>Science</i> 289, 563-4. 4. Katsanis, N. and Fisher, E.M. (1998) <i>Genomics</i> 51, 128-31. 5. Bazirgan, O.A. and Hampton, R.Y. (2008) <i>J Biol Chem</i> 283, 12797-810. 6. Ravid, T. and Hochstrasser, M. (2007) <i>Nat Cell Biol</i> 9, 422-7. 7. Liu, W. et al. (2014) <i>EMBO J</i> 33, 46-61. 8. Li, W. et al. (2007) <i>Nature</i> 446, 333-7. 9. Li, W. et al. (2009) <i>Proc Natl Acad Sci U S A</i> 106, 3722-7. 10. Chen, B. et al. (2006) <i>Proc Natl Acad Sci U S A</i> 103, 341-6.				
Species Reactivi	<b>.</b>	Consider we satisfaction of		g in at least one approve	- d - n- li - ati - n ( - n	

**Species Reactivity** 

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer** 

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X

TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

**Applications Key** 

W: Western Blotting

**Cross-Reactivity Key** 

H: Human M: Mouse R: Rat Mk: Monkey

**Trademarks and Patents** 

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

## **Limited Uses**

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.