

#6338 Store at -20°C

SignalSilence® Merlin siRNA II



✓ 10 µM in 300 µl (100 transfections)

Orders ■ 877-616-CELL (2355) orders@cellsignal.com
Support ■ 877-678-TECH (8324) info@cellsignal.com
Web ■ www.cellsignal.com

rev. 02/10/16

For Research Use Only. Not For Use In Diagnostic Procedures.


Species Cross-Reactivity: H

Description: SignalSilence® Merlin siRNA II from Cell Signaling Technology (CST) allows the researcher to specifically inhibit Merlin expression using RNA interference, a method whereby gene expression can be selectively silenced through the delivery of double stranded RNA molecules into the cell. All SignalSilence® siRNA products from CST are rigorously tested in-house and have been shown to reduce target protein expression by western analysis.

Background: Neurofibromatosis 2 (NF2) is an autosomal dominant inherited disorder that is characterized by the occurrence of vestibular schwannomas, meningiomas and other nervous system tumors. Both the familial tumors of NF2 and equivalent sporadic tumors in the general population are caused by inactivation of the NF2 tumor suppressor gene. Merlin (moesin, ezrin and radixin like protein), the NF2 gene product, has striking similarity to ezrin, radixin and moesin (ERM). Regulation of merlin (aka schwannomin) and ERM proteins involves intramolecular and intermolecular head-to-tail associations between family members (1). Merlin and ERM proteins act as linkers between the plasma membrane and the cytoskeleton, affecting cell morphology, polarity and signal transduction (2). Merlin is phosphorylated by the Rac/Cdc42 effector p21-activated kinase (PAK) at Ser518, negatively regulating Rac (3,4).

Directions for Use: CST recommends transfection with 100 nM Merlin siRNA II 48 to 72 hours prior to cell lysis. For transfection procedure, follow protocol provided by the transfection reagent manufacturer. Please feel free to contact CST with any questions on use.

Quality Control: Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid phase extraction. The annealed RNA duplex is further analyzed by mass spectrometry to verify the exact composition of the duplex. Each lot is compared to the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.



Western blot analysis of extracts from PC3 cells, transfected with 100 nM SignalSilence® Control siRNA (Unconjugated) #6568 (-), SignalSilence® Merlin siRNA I #6316 (+) or SignalSilence® Merlin siRNA II (+), using Phospho-Merlin (Ser518) Antibody #9163 (upper) or α-Tubulin (11H10) Rabbit mAb #2125 (lower). The Phospho-Merlin (Ser518) Antibody confirms silencing of Merlin expression, while the α-Tubulin (11H10) Rabbit mAb is used as a loading control.

Entrez-Gene ID #4771
Swiss-Prot Acc. #P35240

Storage: Merlin siRNA II is supplied in RNase-free water. Aliquot and store at -20°C.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

Background References:

- (1) Ramesh, V. (2004) *Nat. Rev. Neurosci.* 5, 462-470.
- (2) Bretscher, A. et al. (2002) *Nat. Rev. Mol. Cell Biol.* 3, 586-599.
- (3) Xiao, G. H. et al. (2002) *J. Biol. Chem.* 277, 883-886.
- (4) Kissil, J. L. et al. (2003) *Mol. Cell* 12, 841-849.

© 2010 Cell Signaling Technology, Inc.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.