

**MLL2/KMT2B (D6X2E) Rabbit mAb  
(Carboxy-terminal Antigen)**

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP, IF-IC, C&R	H	Endogenous	80	Rabbit IgG	#Q9UMN6	9757

**Product Usage Information**

The CUT&RUN dilution was determined using CUT&RUN Assay Kit #86652.

**Application**

Western Blotting  
Immunoprecipitation  
Immunofluorescence (Immunocytochemistry)  
CUT&RUN

**Dilution**

1:1000  
1:50  
1:1000  
1:50

**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

**Specificity/Sensitivity**

MLL2/KMT2B (D6X2E) Rabbit mAb (Carboxy-terminal Antigen) recognizes endogenous levels of total MLL2 protein. This antibody detects the Taspase 1-cleaved 80 kDa C-terminal MLL2/KMT2B protein (MLL2/KMT2B-C) and the full-length 400 kDa MLL2/KMT2B protein.

**Source / Purification**

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human MLL2/KMT2B protein.

**Background**

The Set1 histone methyltransferase protein was first identified in yeast as part of the Set1/COMPASS histone methyltransferase complex, which methylates histone H3 at Lys4 and functions as a transcriptional co-activator (1). While yeast contain only one known Set1 protein, mammals contain six Set1-related proteins: SET1A, SET1B, MLL1, MLL2, MLL3, and MLL4, all of which assemble into COMPASS-like complexes and methylate histone H3 at Lys4 (2,3). These Set1-related proteins are each found in distinct protein complexes, all of which share the common subunits WDR5, RBBP5, ASH2L, CXXC1, and DPY30, which are required for proper complex assembly and modulation of histone methyltransferase activity (2-6). MLL1 and MLL2 complexes contain the additional protein subunit, menin (6). MLL2, also known as histone-lysine N-methyltransferase 2B (KMT2B), functions to activate gene expression by mediating tri-methylation of histone H3 lysine 4 at the promoters of genes involved in embryogenesis and hematopoiesis, and is required for histone H3 lysine 4 tri-methylation at bivalent promoters in embryonic stem cells (7). Like MLL1, MLL2 is a large protein made up of approximately 2,700 amino acids that is cleaved by the Taspase 1 threonine endopeptidase to form N-terminal (MLL2-N) and C-terminal (MLL2-C) fragments, both of which are subunits of the functional MLL2/COMPASS complex. MLL2-N, MLL2-C, WDR5, RBBP5, and ASH2L define the core catalytic component of the MLL2/COMPASS complex, which is recruited to target genes to regulate transcription. MLL1 gene translocations are often associated with various hematological malignancies and thought to be a driving component of these types of leukemia. MLL2 is required for memory formation, proper glucose homeostasis, and cardiac lineage differentiation of mouse embryonic stem cells (8-11). A recent study has shown that MLL2 is required for survival of MLL-AF9-transformed cells, implicating MLL2 as a potential modulator of MLL1-rearranged leukemias (12). Mutations in MLL2 cause complex early-onset dystonia, and overexpression of MLL2 is associated with gastrointestinal diffuse large B-cell lymphoma (13,14).

**Background References**

1. Miller, T. et al. (2001) *Proc Natl Acad Sci U S A* 98, 12902-7.
2. Shilatifard, A. (2008) *Curr Opin Cell Biol* 20, 341-8.
3. Tenney, K. and Shilatifard, A. (2005) *J Cell Biochem* 95, 429-36.
4. Lee, J.H. and Skalnik, D.G. (2005) *J Biol Chem* 280, 41725-31.
5. Lee, J.H. et al. (2007) *J Biol Chem* 282, 13419-28.
6. Hughes, C.M. et al. (2004) *Mol Cell* 13, 587-97.
7. Denisov, S. et al. (2014) *Development* 141, 526-37.
8. Kerimoglu, C. et al. (2017) *Cell Rep* 20, 538-48.
9. Kerimoglu, C. et al. (2013) *J Neurosci* 33, 3452-64.
10. Goldsworthy, M. et al. (2013) *PLoS One* 8, e61870.
11. Wan, X. et al. (2014) *Stem Cell Rev* 10, 643-52.
12. Chen, Y. et al. (2017) *Cancer Cell* 31, 755-770.e6.
13. Meyer, E. et al. (2017) *Nat Genet* 49, 223-37.

<b>Species Reactivity</b>	Species reactivity is determined by testing in at least one approved application (e.g., western blot).
<b>Western Blot Buffer</b>	<b>IMPORTANT:</b> For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.
<b>Applications Key</b>	<b>W:</b> Western Blotting <b>IP:</b> Immunoprecipitation <b>IF-IC:</b> Immunofluorescence (Immunocytochemistry) <b>C&amp;R:</b> CUT&RUN
<b>Cross-Reactivity Key</b>	<b>H:</b> Human
<b>Trademarks and Patents</b>	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. XP is a registered trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit <a href="http://cellsignal.com/trademarks">cellsignal.com/trademarks</a> for more information.
<b>Limited Uses</b>	<p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p> <p>Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.</p>