

#6564 Store at -20°C

SignalSilence® p38 MAPK siRNA



✓ 10 µM in 300 µl (100 transfections)

Orders ■ 877-616-CELL (2355) orders@cellsignal.com
Support ■ 877-678-TECH (8324) info@cellsignal.com
Web ■ www.cellsignal.com

rev. 03/21/17

For Research Use Only. Not For Use In Diagnostic Procedures.


Species Cross-Reactivity: H

Description: SignalSilence® p38 MAP Kinase siRNA from Cell Signaling Technology (CST) allows the researcher to specifically inhibit p38 MAP Kinase expression using RNA interference, a method whereby gene expression can be selectively silenced through the delivery of double stranded RNA molecules into the cell. All SignalSilence® siRNA products from CST are rigorously tested in-house and have been shown to reduce protein expression by western analysis.

Background: p38 MAP kinase (MAPK), also called RK (1) or CSBP (2), is the mammalian orthologue of the yeast HOG kinase which participates in a signaling cascade controlling cellular responses to cytokines and stress (1-4). Four isoforms of p38 MAP kinase, p38α, β, γ (also known as ERK6 or SAPK3) and δ (also known as SAPK4) have been identified. Similar to the SAPK/JNK pathway, p38 MAP kinase is activated by a variety of cellular stresses including osmotic shock, inflammatory cytokines, lipopolysaccharides (LPS), UV light and growth factors (1-5). MKK3, MKK6 and SEK activate p38 MAP kinase by phosphorylation at Thr180 and Tyr182. Activated p38 MAP kinase has been shown to phosphorylate and activate MAPKAP kinase 2 (3) and to phosphorylate the transcription factors ATF-2 (5), Max (6) and MEF2 (5-8).

Directions for Use: CST recommends transfecting with 100 nM p38 siRNA 48 to 72 hours before lysing cells. For transfection procedure, follow protocol provided by the transfection reagent manufacturer. Please feel free to contact CST with any questions on use.

Quality Control: Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid phase extraction. The annealed RNA duplex is further analyzed by mass spectrometry to verify the exact composition of the duplex. Each lot is compared to the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.



Western blot analysis of extracts from HeLa cells, transfected with 100 nM SignalSilence® Control siRNA (Unconjugated) #6568 (-), SignalSilence® p38 MAPK siRNA I (+) or SignalSilence® p38 MAPK siRNA II #6243 (+), using p38 MAPK Antibody #9212 and α-Tubulin (11H10) Rabbit mAb #2125. The p38 MAPK antibody confirms silencing of p38 MAPK expression while the α-Tubulin (11H10) Rabbit mAb is used to control for loading and specificity of p38 MAPK siRNA.

Entrez-Gene ID #1432
Swiss-Prot Acc. #Q16539

Storage: p38 MAPK siRNA I is supplied in RNase-free water. Aliquot and store at -20°C.

Please visit www.cellsignal.com for a complete listing of recommended companion products.

Background References:

- (1) Rouse, J. et al. (1994) *Cell* 78, 1027-1037.
- (2) Han, J. et al. (1994) *Science* 265, 808-811.
- (3) Lee, J.C. et al. (1994) *Nature* 372, 739-746.
- (4) Freshney, N.W. et al. (1994) *Cell* 78, 1039-1049.
- (5) Raingeaud, J. et al. (1995) *J. Biol. Chem.* 270, 7420-7426.
- (6) Zervos, A.S. et al. (1995) *Proc. Natl. Acad. Sci. USA* 92, 10531-10534.
- (7) Zhao, M. et al. (1999) *Mol. Cell. Biol.* 19, 21-30.
- (8) Yang, S.H. et al. (1999) *Mol. Cell. Biol.* 19, 4028-4038.

© 2009 Cell Signaling Technology, Inc.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.