

ARID1B/BAF250B (E1U7D) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP, IHC-P	H M	Endogenous	250, 280	Rabbit IgG	#Q8NFD5	57492

Product Usage Information**Application**

Western Blotting
Immunoprecipitation
Immunohistochemistry (Paraffin)

Dilution

1:1000
1:100
1:500

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

ARID1B/BAF250B (E1U7D) Rabbit mAb recognizes endogenous levels of total ARID1B/BAF250B protein. This antibody does not cross-react with ARID1A/BAF250A protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Ala1320 of human ARID1B/BAF250B protein.

Background

ATP-dependent chromatin remodeling complexes play an essential role in the regulation of nuclear processes such as transcription and DNA replication and repair (1,2). The SWI/SNF chromatin remodeling complex consists of more than 10 subunits and contains a single molecule of either BRM or BRG1 as the ATPase catalytic subunit. The activity of the ATPase subunit disrupts histone-DNA contacts and changes the accessibility of crucial regulatory elements to the chromatin. The additional core and accessory subunits play a scaffolding role to maintain stability and provide surfaces for interaction with various transcription factors and chromatin (2-5). The interactions between SWI/SNF subunits and transcription factors, such as nuclear receptors, p53, Rb, BRCA1, and MyoD, facilitate recruitment of the complex to target genes for regulation of gene activation, cell growth, cell cycle, and differentiation processes (1,6-9).

ARID1B (A-T rich interacting domain 1B), also known as BAF250B, is a DNA-binding member of the SWI/SNF complex. It has 60% homology with ARID1A/BAF250A, and the proteins are mutually exclusive members of the complex, akin to Brg1 and BRM (10). ARID1B plays a role in synapse formation and dendritic arborization in neuronal development, and haploinsufficiency of ARID1B has been reported in intellectual disability (11-13). Mutations in ARID1B have also been shown in Coffin-Siris syndrome (14). ARID1B/BAF250B is a critical vulnerability in ARID1A/BAF250A mutant cancers, and could be explored as a potential therapeutic target (15).

Background References

1. Ho, L. and Crabtree, G.R. (2010) *Nature* 463, 474-84.
2. Becker, P.B. and Hörz, W. (2002) *Annu Rev Biochem* 71, 247-73.
3. Eberharder, A. and Becker, P.B. (2004) *J Cell Sci* 117, 3707-11.
4. Bowman, G.D. (2010) *Curr Opin Struct Biol* 20, 73-81.
5. Gangaraju, V.K. and Bartholomew, B. (2007) *Mutat Res* 618, 3-17.
6. Lessard, J.A. and Crabtree, G.R. (2010) *Annu Rev Cell Dev Biol* 26, 503-32.
7. Morettini, S. et al. (2008) *Front Biosci* 13, 5522-32.
8. Wolf, I.M. et al. (2008) *J Cell Biochem* 104, 1580-6.
9. Simone, C. (2006) *J Cell Physiol* 207, 309-14.
10. Wang, X. et al. (2004) *Biochem J* 383, 319-25.
11. Ka, M. et al. (2016) *J Neurosci* 36, 2723-42.
12. Halgren, C. et al. (2012) *Clin Genet* 82, 248-55.
13. Hoyer, J. et al. (2012) *Am J Hum Genet* 90, 565-72.
14. Tsurusaki, Y. et al. (2012) *Nat Genet* 44, 376-8.
15. Helming, K.C. et al. (2014) *Nat Med* 20, 251-4.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation **IHC-P:** Immunohistochemistry (Paraffin)

Cross-Reactivity Key

H: Human **M:** Mouse

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

SignalStain is a registered trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.