

TBC1D1 (D2Y8M) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications: W, IP	Reactivity: M	Sensitivity: Endogenous	MW (kDa): 160	Source/Isotype: Rabbit IgG	UniProt ID: #Q60949	Entrez-Gene Id: 57915
-------------------------------	-------------------------	-----------------------------------	-------------------------	--------------------------------------	-------------------------------	---------------------------------

Product Usage Information	Application Western Blotting Immunoprecipitation	Dilution 1:1000 1:50
Storage	Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.	
Specificity/Sensitivity	TBC1D1 (D2Y8M) Rabbit mAb recognizes endogenous levels of total TBC1D1 protein.	
Source / Purification	Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Val796 of mouse TBC1D1 protein.	
Background	TBC1D1 is a paralog of AS160 (1) and both proteins share about 50% identity (2). TBC1D1 was shown to be a candidate gene for severe obesity (3). It plays a role in Glut4 translocation through its GAP activity (2,4). Studies indicate that TBC1D1 is highly expressed in skeletal muscle (1). Insulin, AICAR, and contraction directly regulate TBC1D1 phosphorylation in this tissue (1). Three AMPK phosphorylation sites (Ser231, Ser660, and Ser700) and one Akt phosphorylation site (Thr590) were identified in skeletal muscle (5). Muscle contraction or AICAR treatment increases phosphorylation on Ser231, Ser660, and Ser700 but not on Thr590; insulin increases phosphorylation on Thr590 only (5).	
Background References	<ol style="list-style-type: none"> 1. Taylor, E.B. et al. (2008) <i>J Biol Chem</i> 283, 9787-96. 2. Roach, W.G. et al. (2007) <i>Biochem J</i> 403, 353-8. 3. Stone, S. et al. (2006) <i>Hum Mol Genet</i> 15, 2709-20. 4. Chavez, J.A. et al. (2008) <i>J Biol Chem</i> 283, 9187-95. 5. Vichaiwong, K. et al. (2010) <i>Biochem J</i> 431, 311-20. 	
Species Reactivity	Species reactivity is determined by testing in at least one approved application (e.g., western blot).	
Western Blot Buffer	IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.	
Applications Key	W: Western Blotting IP: Immunoprecipitation	
Cross-Reactivity Key	M: Mouse	
Trademarks and Patents	Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc. All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.	
Limited Uses	<p>Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.</p> <p>Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not</p>	

sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.