CD161/KLRB1 (HP-3G10) Mouse mAb (FITC Conjugate)



Orders: 877-616-CELL (2355) orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com

cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

preparation. This Cell Signaling Technology antibody is conjugated to FITC and tested in-house for direct flow cytometric analysis in human cells. Background CD161/KLRB1 (Killer cell lectin-like receptor subfamily B member 1, also known as CLEC5B and NKR-P1A) is a type II transmembrane protein that is expressed on the majority of Natural Killer (NK) cells, T cells, and some T lymphocytes (1). CD161/KLRB1 is also expressed on Th17 cells, promotes their generation, and modulates their function (2). Engagement with its ligand lectin-like transcript 1 (LLT1 inhibits NK cell function, while LLT1 and CD161/KLRB1 interaction in the presence of a TCR signal enhances IFN-gamma production by T cells (3,4). There are several different CD161 isoforms in roden and some function as activating receptors as well (5,6). This HP-3G10 antibody is used for flow cytometric analysis of CD161/KLRB1 expression on NK, NKT, a various subsets of T cells. Background References 1. Lanier, L.L. et al. (1994) / Immunol 153, 2417-28. 2. Bai, A. et al. (2014) / Immunol 193, 3366-77.	Applications: FC-L	Reactivity:	Sensitivity: Endogenous	Source/Isotype: Mouse IgG1 kappa	UniProt ID: #Q12918	Entrez-Gene Id: 3820
for 12 months when stored at 4°C. Do not aliquot the antibody. Protect from light. Do not freeze. Specificity/Sensitivity CD161/KLRB1 (HP-3G10) Mouse mAb (FITC Conjugate) recognizes endogenous levels of total CD161/KLRB1 protein. This antibody detects an epitope within the extracellular domain. Source / Purification This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. Description This Cell Signaling Technology antibody is conjugated to FITC and tested in-house for direct flow cytometric analysis in human cells. Background CD161/KLRB1 (Killer cell lectin-like receptor subfamily B member 1, also known as CLEC5B and NKR-P1A) is a type II transmembrane protein that is expressed on the majority of Natural Killer (NK) cells, I cells, and some T lymphocytes (1). CD161/KLRB1 is also expressed on Th17 cells, promotes their generation, and modulates their function (2). Engagement with its ligand lectin-like transcript 1 (LLT1 inhibits NK cell function, while LLT1 and CD161/KLRB1 interaction in the presence of a TCR signal enhances IFN-gamma production by T cells (3,4). There are several different CD161 isoforms in roden and some function as activating receptors as well (5,6). This HP-3G10 antibody is used for flow cytometric analysis of CD161/KLRB1 expression on NK, NKT, a various subsets of T cells. Background References 1. Lanier, L.L. et al. (1994) J Immunol 153, 2417-28. 2. Bal, A. et al. (2014) J Immunol 193, 3366-77.	Product Usage Information		• • • • • • • • • • • • • • • • • • • •			
CD161/KLRB1 protein. This antibody detects an epitope within the extracellular domain. Source / Purification This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. Description This Cell Signaling Technology antibody is conjugated to FITC and tested in-house for direct flow cytometric analysis in human cells. Background CD161/KLRB1 (Killer cell lectin-like receptor subfamily B member 1, also known as CLEC5B and NKR-P1A) is a type II transmembrane protein that is expressed on the majority of Natural Killer (NK) cells, T cells, and some T lymphocytes (1). CD161/KLRB1 is also expressed on Th17 cells, promotes their generation, and modulates their function (2). Engagement with its ligand lectin-like transcript 1 (LLT1 inhibits NK cell function, while LLT1 and CD161/KLRB1 interaction in the presence of a TCR signal enhances IFN-gamma production by T cells (3,4). There are several different CD161 isoforms in roden and some function as activating receptors as well (5,6). This HP-3G10 antibody is used for flow cytometric analysis of CD161/KLRB1 expression on NK, NKT, a various subsets of T cells. Background References 1. Lanier, L.L. et al. (1994) J Immunol 153, 2417-28. 2. Bai, A. et al. (2014) J Immunol 193, 3366-77.	Storage					
The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. This Cell Signaling Technology antibody is conjugated to FITC and tested in-house for direct flow cytometric analysis in human cells. Background CD161/KLRB1 (Killer cell lectin-like receptor subfamily B member 1, also known as CLEC5B and NKR-P1A) is a type II transmembrane protein that is expressed on the majority of Natural Killer (NK) cells, in tells, and some T lymphocytes (1). CD161/KLRB1 is also expressed on Th17 cells, promotes their generation, and modulates their function (2). Engagement with its ligand lectin-like transcript 1 (LLT1 inhibits NK cell function, while LLT1 and CD161/KLRB1 interaction in the presence of a TCR signal enhances IFN-gamma production by T cells (3,4). There are several different CD161 isoforms in roden and some function as activating receptors as well (5,6). This HP-3G10 antibody is used for flow cytometric analysis of CD161/KLRB1 expression on NK, NKT, a various subsets of T cells. Background References 1. Lanier, L.L. et al. (1994) J Immunol 153, 2417-28. 2. Bai, A. et al. (2014) J Immunol 193, 3366-77.	Specificity/Sensitivity					
cytometric analysis in human cells. Background CD161/KLRB1 (Killer cell lectin-like receptor subfamily B member 1, also known as CLEC5B and NKR-P1A) is a type II transmembrane protein that is expressed on the majority of Natural Killer (NK) cells, I cells, and some T lymphocytes (1). CD161/KLRB1 is also expressed on Th17 cells, promotes their generation, and modulates their function (2). Engagement with its ligand lectin-like transcript 1 (LLT1 inhibits NK cell function, while LLT1 and CD161/KLRB1 interaction in the presence of a TCR signal enhances IFN-gamma production by T cells (3,4). There are several different CD161 isoforms in roden and some function as activating receptors as well (5,6). This HP-3G10 antibody is used for flow cytometric analysis of CD161/KLRB1 expression on NK, NKT, a various subsets of T cells. Background References 1. Lanier, L.L. et al. (1994) J Immunol 153, 2417-28. 2. Bai, A. et al. (2014) J Immunol 193, 3366-77.	Source / Purification		The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the			
P1A) is a type II transmembrane protein that is expressed on the majority of Natural Killer (NK) cells, I T cells, and some T lymphocytes (1). CD161/KLRB1 is also expressed on Th17 cells, promotes their generation, and modulates their function (2). Engagement with its ligand lectin-like transcript 1 (LLT1 inhibits NK cell function, while LLT1 and CD161/KLRB1 interaction in the presence of a TCR signal enhances IFN-gamma production by T cells (3,4). There are several different CD161 isoforms in roden and some function as activating receptors as well (5,6). This HP-3G10 antibody is used for flow cytometric analysis of CD161/KLRB1 expression on NK, NKT, a various subsets of T cells. Background References 1. Lanier, L.L. et al. (1994) J Immunol 153, 2417-28. 2. Bai, A. et al. (2014) J Immunol 193, 3366-77.	Description		5 5 5 7 7 5			
2. Bai, A. et al. (2014) <i>J Immunol</i> 193, 3366-77.	Background		P1A) is a type II transmembrane protein that is expressed on the majority of Natural Killer (NK) cells, NK T cells, and some T lymphocytes (1). CD161/KLRB1 is also expressed on Th17 cells, promotes their generation, and modulates their function (2). Engagement with its ligand lectin-like transcript 1 (LLT1) inhibits NK cell function, while LLT1 and CD161/KLRB1 interaction in the presence of a TCR signal enhances IFN-gamma production by T cells (3,4). There are several different CD161 isoforms in rodents and some function as activating receptors as well (5,6). This HP-3G10 antibody is used for flow cytometric analysis of CD161/KLRB1 expression on NK, NKT, and			
4. Rosen, D.B. et al. (2005) <i>J Immunol</i> 175, 7796-9. 5. Carlyle, J.R. et al. (2006) <i>J Immunol</i> 176, 7511-24. 6. Kirkham, C.L. and Carlyle, J.R. (2014) <i>Front Immunol</i> 5, 214.	Background References		2. Bai, A. et al. (2014) <i>J Immunol</i> 193, 3366-77. 3. Aldemir, H. et al. (2005) <i>J Immunol</i> 175, 7791-5. 4. Rosen, D.B. et al. (2005) <i>J Immunol</i> 175, 7796-9. 5. Carlyle, J.R. et al. (2006) <i>J Immunol</i> 176, 7511-24.			

Species Reactivity Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Applications Key FC-L: Flow Cytometry (Live)

Cross-Reactivity Key H: Human

Trademarks and Patents Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for

more information.

Limited UsesExcept as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's

terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no

force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a

component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com For Research Use Only. Not for Use in Diagnostic Procedures.