

Store at  
-20C  
#68309

# Notch Activated Targets Antibody Sampler Kit



Orders: 877-616-CELL (2355)  
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com  
cellsignal.com

1 Kit (8 x 20 microliters)

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Quantity	Mol. Wt	Isotype/Source
Notch1 (D1E11) XP <sup>®</sup> Rabbit mAb	3608	20 µl	120, 300 kDa	Rabbit IgG
Cleaved Notch1 (Val1744) (D3B8) Rabbit mAb	4147	20 µl	110 kDa	Rabbit IgG
RBPSUH (D10A4) XP <sup>®</sup> Rabbit mAb	5313	20 µl	61 kDa	Rabbit IgG
MAML1 (D3K7B) Rabbit mAb	12166	20 µl	130 kDa	Rabbit IgG
c-Myc (D84C12) Rabbit mAb	5605	20 µl	57-65 kDa	Rabbit IgG
p21 Waf1/Cip1 (12D1) Rabbit mAb	2947	20 µl	21 kDa	Rabbit IgG
HES1 (D6P2U) Rabbit mAb	11988	20 µl	30 kDa	Rabbit IgG
Cyclin D3 (DCS22) Mouse mAb	2936	20 µl	31 kDa	Mouse IgG1
Anti-rabbit IgG, HRP-linked Antibody	7074	100 µl		Goat
Anti-mouse IgG, HRP-linked Antibody	7076	100 µl		Horse

Please visit [cellsignal.com](http://cellsignal.com) for individual component applications, species cross-reactivity, dilutions, protocols, and additional product information.

## Description

The Notch Activated Targets Antibody Sampler Kit provides an economical means of detecting target proteins of activated Notch. The kit contains enough primary antibody to perform four western blot experiments per primary antibody.

## Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

## Background

Notch proteins (Notch1-4) are a family of transmembrane receptors that play important roles in development and the determination of cell fate (1). Mature Notch receptors are processed and assembled as heterodimeric proteins, with each dimer comprised of a large extracellular ligand-binding domain, a single-pass transmembrane domain, and a smaller cytoplasmic subunit (Notch intracellular domain, NICD) (2). Binding of Notch receptors to ligands of the Delta-Serrate-Lag2 (DSL) family triggers heterodimer dissociation, exposing the receptors to proteolytic cleavages; these result in release of the NICD, which translocates to the nucleus and activates transcription of downstream target genes (3,4). RBPSUH (Recombining Binding Protein, SUppressor of Hairless), is the DNA-binding component of the transcription complex regulated by canonical Notch signaling. Binding of Notch with RBPSUH activates a transcription activation complex that includes Mastermind-like (MAML) proteins, leading to transcriptional activation of Notch target genes (5-7). The NICD binds and activates c-Myc which functions as a transcriptional regulator with roles in various aspects of cell behavior including proliferation, differentiation and apoptosis (8). The tumor suppressor protein p21 Waf1/Cip1 acts as an inhibitor of cell cycle progression. The NICD-RBPSUH complex binds and activates p21 for transcription (15). HES1 (Hairy and Enhancer of Split 1) is one of seven members of the HES family of basic helix-loop-helix (bHLH) transcription factors that is particularly well known as a repressive mediator of the canonical Notch signaling pathway (10). HES1 plays a key role in mediating Notch-dependent T cell lineage commitment (11), and has been reported to be an essential mediator of Notch-induced T cell acute lymphoblastic leukemia (T-ALL) (11,12). The active complex of cyclin D/CDK4 targets the retinoblastoma protein for phosphorylation, allowing the release of E2F transcription factors that activate G1/S-phase gene expression (13). Transcription of cyclin D is in part regulated by the NICD binding to the promoter region of cyclin D (14).

## Background References

1. Artavanis-Tsakonas, S. et al. (1999) *Science* 284, 770-6.
2. Chan, Y.M. and Jan, Y.N. (1998) *Cell* 94, 423-6.
3. Schroeter, E.H. et al. (1998) *Nature* 393, 382-6.
4. Rand, M.D. et al. (2000) *Mol Cell Biol* 20, 1825-35.
5. Wu, L. et al. (2002) *Mol Cell Biol* 22, 7688-700.
6. Lin, S.E. et al. (2002) *J Biol Chem* 277, 50612-20.
7. Kitagawa, M. et al. (2001) *Mol Cell Biol* 21, 4337-46.
8. Baudino, T.A. and Cleveland, J.L. (2001) *Mol Cell Biol* 21, 691-702.

9. Flores-Rozas, H. et al. (1994) *Proc Natl Acad Sci U S A* 91, 8655-9.
  10. Kobayashi, T. and Kageyama, R. (2010) *Genes Cells* 15, 689-98.
  11. Wendorff, A.A. et al. (2010) *Immunity* 33, 671-84.
  12. Espinosa, L. et al. (2010) *Cancer Cell* 18, 268-81.
  13. Lukas, J. et al. (1996) *Mol Cell Biol* 16, 6917-25.
  14. Li, X. and von Boehmer, H. (2011) *ISRN Hematol* 2011, 921706.
  15. Niimi, H. et al. (2007) *J Cell Biol* 176, 695-707.
- 

## Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.