

Store at
-20C
#68348**GIRK2 (D3T1N) Rabbit mAb**
Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W, IP	H M R	Endogenous	46	Rabbit IgG	#P48051	3763

Product Usage Information**Application**

Western Blotting
Immunoprecipitation

Dilution

1:1000
1:50

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity
Source / Purification

GIRK2 (D3T1N) Rabbit mAb recognizes endogenous levels of total GIRK2 protein.

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Glu22 of human GIRK2 protein.

Background

GIRK2 is a member of G protein-coupled inwardly rectifying potassium channel family proteins (GIRKs). GIRK family proteins allow potassium to flow into the cell and therefore control cellular excitability in the central nervous system, heart, and pancreas (1-4). Activation of most GIRK channels requires heterologous subunit assembly and the presence of ATP (5-7). GIRK2 is abundantly expressed in the brain, where it is involved in pain perception. It is also required for peripheral opioid-mediated analgesia (8). Additionally GIRK2 localizes to pancreatic β cells and regulates insulin secretion (9,10). Mutations in the *KCNJ6* gene encoding GIRK2 are associated with Keppen-Lubinsky Syndrome, a rare disease characterized by severe developmental delay, facial dysmorphism, and intellectual disability (11).

Background References

1. Lesage, F. et al. (1995) *J Biol Chem* 270, 28660-7.
2. Kubo, Y. et al. (1993) *Nature* 364, 802-6.
3. Krapivinsky, G. et al. (1995) *Nature* 374, 135-41.
4. Isomoto, S. et al. (1997) *Jpn J Physiol* 47, 11-39.
5. Duprat, F. et al. (1995) *Biochem Biophys Res Commun* 212, 657-63.
6. Corey, S. and Clapham, D.E. (1998) *J Biol Chem* 273, 27499-504.
7. Han, J. et al. (2003) *J Physiol* 550, 693-706.
8. Nockemann, D. et al. (2013) *EMBO Mol Med* 5, 1263-77.
9. Ferrer, J. et al. (1995) *J Biol Chem* 270, 26086-91.
10. Sakura, H. et al. (1995) *FEBS Lett* 367, 193-7.
11. Masotti, A. et al. (2015) *Am J Hum Genet* 96, 295-300.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting **IP:** Immunoprecipitation

Cross-Reactivity Key

H: Human **M:** Mouse **R:** Rat

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

XP is a registered trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.