

Store at
-20C
#69493**CD177 (F905U) & CO-0202-488 SignalStar®
Oligo-Antibody Pair****Orders:** 877-616-CELL (2355)
orders@cellsignal.com**Support:** 877-678-TECH (8324)**Web:** info@cellsignal.com
cellsignal.com

1 Kit (10 slides)

UniProt ID: #Q8N6Q3
Entrez-Gene Id: #57126

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Product Includes	Product #	Volume	Reactivity	Isotype
CD177 (F905U) Rabbit Monoclonal Antibody (SignalStar™ Conjugate 0202)	78364	50 µl	H	Rabbit IgG
Complementary Oligo (CO-0202-488)	86824	30 µl		

Storage

SignalStar conjugates are supplied in PBS (pH 7.2), less than 0.1% sodium azide, 2 mM EDTA, 0.05% Triton X-100, 2 mg/mL BSA, and 50% glycerol. Complementary oligos are supplied in nuclease-free water. Store at -20°C. *Do not aliquot the antibody.* All components in this kit are stable for at least 12 months when stored at the recommended temperature.

Description

SignalStar multiplex immunohistochemistry (IHC) is an advanced technology for labeling multiple proteins simultaneously in tissue samples using specific primary antibodies and fluorescent detection reagents. This technology offers accuracy and reliability in visualizing and analyzing protein expression while maintaining spatial context and tissue architecture.

SignalStar Oligo-Antibody Pairs are compatible with the SignalStar Multiplex IHC Buffer Kits for use in fluorescent multiplex imaging experiments. This product includes the oligo-conjugated antibodies and complementary oligos required for labeling your target protein on up to 10 slides. SignalStar Multiplex IHC Buffer Kits are required to amplify and image the target signal. Multiple oligo-antibody pairs can be conveniently combined into a multiplex panel using the SignalStar Multiplex IHC Panel Builder. SignalStar Multiplex IHC Kits & Reagents are not compatible with all of Cell Signaling Technology® products and protocols that are recommended for use in immunohistochemical assays.

Specificity/Sensitivity

CD177 (F905U) Rabbit Monoclonal Antibody (SignalStar™ Conjugate 0202) recognizes endogenous levels of total CD177 protein. Non-specific staining was observed in human normal and neoplastic kidney by immunohistochemistry.

Source / Purification

Monoclonal antibody is produced by immunizing animals with recombinant protein specific to the amino terminus of human CD177 protein.

Background

Cluster of differentiation 177 antigen (CD177, also known as human neutrophil alloantigen 2a and NB1), is a glycosyl-phosphatidylinositol (GPI)-linked N-glycosylated transmembrane protein expressed primarily on the surface of neutrophils (1,2). Expression of CD177 can be induced by granulocyte-colony-stimulating factor and during pregnancy or bacterial infection (3-5). CD177 mediates adhesion and transendothelial migration of neutrophils via interaction with CD31/PECAM-1 and integrin β2 on endothelial cells (6,7). CD177 also promotes the cell surface expression of proteinase 3 (PR3), which may play a role in enhancing endothelial cell junctional integrity and thus vascular integrity during neutrophil diapedesis (8-10). In association with CD11b/ITGAM, integrin β2, and PR3, CD177 promotes activation of TNF-α primed neutrophils including degranulation and superoxide production (11). CD177 is a polymorphic gene that has been linked to several important clinical diseases, including polycythemia vera, Wegener's granulomatosis, and immune mediated neonatal neutropenia (12-14).

Background References

1. Matsuo, K. et al. (2000) *Transfusion* 40, 654-62.
2. Xie, Q. et al. (2015) *Protein Cell* 6, 117-26.
3. Stroncek, D.F. et al. (1998) *Transfusion* 38, 663-8.
4. Caruccio, L. et al. (2003) *Transfusion* 43, 357-63.
5. Göhring, K. et al. (2004) *Br J Haematol* 126, 252-4.
6. Sachs, U.J. et al. (2007) *J Biol Chem* 282, 23603-12.
7. Bai, M. et al. (2017) *Blood* 130, 2092-100.
8. von Vietinghoff, S. et al. (2007) *Blood* 109, 4487-93.
9. von Vietinghoff, S. et al. (2008) *Clin Exp Immunol* 152, 508-16.
10. Kuckleburg, C.J. and Newman, P.J. (2013) *Arterioscler Thromb Vasc Biol* 33, 275-84.
11. Jerke, U. et al. (2011) *J Biol Chem* 286, 7070-81.
12. Lalezari, P. et al. (1971) *J Clin Invest* 50, 1108-15.
13. Bettinotti, M.P. et al. (2002) *Clin Immunol* 102, 138-44.
14. Caruccio, L. et al. (2006) *Transfusion* 46, 441-7.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

SignalStar is a registered trademark of Cell Signaling Technology, Inc.

U.S. Patent No. 10,781,477, foreign equivalents, and child patents deriving therefrom.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.