

**UBE2N/Ubc13 (D2A1) Rabbit mAb**

**Orders:** 877-616-CELL (2355)  
orders@cellsignal.com

**Support:** 877-678-TECH (8324)

**Web:** info@cellsignal.com  
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

**For Research Use Only. Not for Use in Diagnostic Procedures.**

<b>Applications:</b> W, IP	<b>Reactivity:</b> H M R Mk	<b>Sensitivity:</b> Endogenous	<b>MW (kDa):</b> 17	<b>Source/Isotype:</b> Rabbit IgG	<b>UniProt ID:</b> #P61088	<b>Entrez-Gene Id:</b> 7334
-------------------------------	--------------------------------	-----------------------------------	------------------------	--------------------------------------	-------------------------------	--------------------------------

**Product Usage Information****Application**

Western Blotting  
Immunoprecipitation

**Dilution**

1:1000  
1:100

**Storage**

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

**Specificity/Sensitivity**

UBE2N/Ubc13 (D2A1) Rabbit mAb recognizes endogenous levels of total UBE2N/Ubc13 protein. This antibody does not cross-react with UBE2D family members.

**Species predicted to react based on 100% sequence homology**

Chicken, Xenopus, Bovine, Dog, Horse

**Source / Purification**

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human UBE2N/Ubc13 protein.

**Background**

Protein ubiquitination is an important posttranslational modification that regulates protein function and fate (1). Ubiquitin (Ub) can be conjugated to target proteins in either monomeric or polymeric forms. There are several different lysine residues within Ub that can be used as conjugation sites for poly-Ub chain formation. Different poly-Ub linkages mediate different functions of the target protein ranging from alterations in protein function to degradation (2). UBE2N/Ubc13 is a ubiquitin-E2-conjugating enzyme that catalyzes K63-linked poly-Ub chain formation (1,2). UBE2N forms a heterodimer with MMS2 or Uev1A to exert its E2 ligase function. The UBE2N/MMS2 and UBE2N/Uev1A heterodimers catalyze different modes of target protein ubiquitination to mediate various signaling pathways (3-5) including: DNA damage and recombination, p53 and check point control, the cell cycle (6-10), immunoreceptor signaling (11,12), and endocytosis (13). Most recently, UBE2N was shown to play an important role in inflammatory signaling by promoting K63-linked ubiquitination and activation of IKK downstream of the IL-1β receptor (14). Furthermore, interaction of UBE2N with the Triad1 E3 protein-ubiquitin ligase was shown to play an important role in myelopoiesis (15).

**Background References**

- Herrmann, J. et al. (2007) *Circ Res* 100, 1276-91.
- Wilkinson, K.D. et al. (2005) *EMBO Rep* 6, 815-20.
- Hofmann, R.M. and Pickart, C.M. (1999) *Cell* 96, 645-53.
- Deng, L. et al. (2000) *Cell* 103, 351-61.
- Andersen, P.L. et al. (2005) *J Cell Biol* 170, 745-55.
- Zhao, G.Y. et al. (2007) *Mol Cell* 25, 663-75.
- Kolas, N.K. et al. (2007) *Science* 318, 1637-40.
- Laine, A. et al. (2006) *Mol Cell Biol* 26, 8901-13.
- Huen, M.S. et al. (2008) *Mol Cell Biol* 28, 6104-12.
- Loring, G.L. et al. (2008) *Cell Cycle* 7, 96-105.
- Yamamoto, M. et al. (2006) *Nat Immunol* 7, 962-70.
- Yamamoto, M. et al. (2006) *J Immunol* 177, 7520-4.
- Duncan, L.M. et al. (2006) *EMBO J* 25, 1635-45.
- Xu, M. et al. (2009) *Mol Cell* 36, 302-14.
- Marteijn, J.A. et al. (2009) *Leukemia* 23, 1480-9.

**Species Reactivity**

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

**Western Blot Buffer**

**IMPORTANT:** For western blots, incubate membrane with diluted primary antibody in 5% w/v BSA, 1X TBS, 0.1% Tween@ 20 at 4°C with gentle shaking, overnight.

**Applications Key**

**W:** Western Blotting **IP:** Immunoprecipitation

## Cross-Reactivity Key

**H:** Human **M:** Mouse **R:** Rat **Mk:** Monkey

## Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit [cellsignal.com/trademarks](http://cellsignal.com/trademarks) for more information.

## Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.