

P2X4 Receptor (D9R1H) Rabbit mAb

Orders: 877-616-CELL (2355)
orders@cellsignal.com

Support: 877-678-TECH (8324)

Web: info@cellsignal.com
cellsignal.com

3 Trask Lane | Danvers | Massachusetts | 01923 | USA

For Research Use Only. Not for Use in Diagnostic Procedures.

Applications:	Reactivity:	Sensitivity:	MW (kDa):	Source/Isotype:	UniProt ID:	Entrez-Gene Id:
W	H	Endogenous	62	Rabbit IgG	#Q99571	5025

Product Usage Information**Application**

Western Blotting

Dilution

1:1000

Storage

Supplied in 10 mM sodium HEPES (pH 7.5), 150 mM NaCl, 100 µg/ml BSA, 50% glycerol and less than 0.02% sodium azide. Store at -20°C. Do not aliquot the antibody.

Specificity/Sensitivity

P2X4 Receptor (D9R1H) Rabbit mAb recognizes endogenous levels of total P2X4 receptor protein.

Source / Purification

Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues surrounding Pro120 of human P2X4 receptor protein.

Background

P2X purinergic receptors are ATP-gated ion channels involved in physiological processes that include inflammation, afferent sensory signaling, and sympathetic motor nerve activity. Seven different vertebrate genes (*P2RX1-P2RX7*) encode for individual receptor protein subunits (1). All P2X subunit proteins share similar protein domain structure, but can differ in overall protein length from 384 to 595 amino acids. Each P2X subunit is composed of amino- and carboxy-terminal intracellular domains, two transmembrane domains, and a large extracellular loop that contains ten evenly spaced cysteines and multiple glycosylation sites (2). P2X receptors are found in a variety of cell types and tissues, including central and peripheral nervous system neurons and glial cells, autonomic and sensory neurons, bone, muscle, and hematopoietic tissues (1).

P2X purinoceptor 4 (P2X4) trimers are expressed at the cell surface where they act as ligand-gated ion channels for monovalent and divalent cations (3). P2X4 receptors contribute to regulation of synaptic strength through participation in the formation of long-term potentiation (4,5). Research studies indicate that P2X4 receptor expression may play a role in influencing alcohol-drinking behavior and conferring protection to cardiac myocytes during heart failure (6,7). Additional studies show that microglial P2X4 receptors are upregulated during nerve-injury associated neuropathic pain (8).

Background References

1. North, R.A. (2002) *Physiol Rev* 82, 1013-67.
2. Valera, S. et al. (1994) *Nature* 371, 516-9.
3. Abbracchio, M.P. et al. (2009) *Trends Neurosci* 32, 19-29.
4. Fujii, S. (2004) *J Pharmacol Sci* 94, 103-6.
5. Sim, J.A. et al. (2006) *J Neurosci* 26, 9006-9.
6. Franklin, K.M. et al. (2014) *Front Neurosci* 8, 176.
7. Yang, T. et al. (2014) *Circ Heart Fail* 7, 510-8.
8. Tsuda, M. et al. (2013) *Front Cell Neurosci* 7, 191.

Species Reactivity

Species reactivity is determined by testing in at least one approved application (e.g., western blot).

Western Blot Buffer

IMPORTANT: For western blots, incubate membrane with diluted primary antibody in 5% w/v nonfat dry milk, 1X TBS, 0.1% Tween® 20 at 4°C with gentle shaking, overnight.

Applications Key

W: Western Blotting

Cross-Reactivity Key

H: Human

Trademarks and Patents

Cell Signaling Technology is a trademark of Cell Signaling Technology, Inc.

All other trademarks are the property of their respective owners. Visit cellsignal.com/trademarks for more information.

Limited Uses

Except as otherwise expressly agreed in a writing signed by a legally authorized representative of CST, the following terms apply to Products provided by CST, its affiliates or its distributors. Any Customer's terms and conditions that are in addition to, or different from, those contained herein, unless separately accepted in writing by a legally authorized representative of CST, are rejected and are of no force or effect.

Products are labeled with For Research Use Only or a similar labeling statement and have not been approved, cleared, or licensed by the FDA or other regulatory foreign or domestic entity, for any purpose. Customer shall not use any Product for any diagnostic or therapeutic purpose, or otherwise in any manner that conflicts with its labeling statement. Products sold or licensed by CST are provided for Customer as the end-user and solely for research and development uses. Any use of Product for diagnostic, prophylactic or therapeutic purposes, or any purchase of Product for resale (alone or as a component) or other commercial purpose, requires a separate license from CST. Customer shall (a) not sell, license, loan, donate or otherwise transfer or make available any Product to any third party, whether alone or in combination with other materials, or use the Products to manufacture any commercial products, (b) not copy, modify, reverse engineer, decompile, disassemble or otherwise attempt to discover the underlying structure or technology of the Products, or use the Products for the purpose of developing any products or services that would compete with CST products or services, (c) not alter or remove from the Products any trademarks, trade names, logos, patent or copyright notices or markings, (d) use the Products solely in accordance with CST Product Terms of Sale and any applicable documentation, and (e) comply with any license, terms of service or similar agreement with respect to any third party products or services used by Customer in connection with the Products.

Orders: 877-616-CELL (2355) • orders@cellsignal.com • Support: 877-678-TECH (8324) • info@cellsignal.com • Web: cellsignal.com
For Research Use Only. Not for Use in Diagnostic Procedures.